

# Product Datasheet

## CXCL8/IL-8 Antibody - BSA Free NBP2-33819

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

### Publications: 5

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-33819](http://www.novusbio.com/NBP2-33819)

Updated 2/23/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-33819](http://www.novusbio.com/reviews/destination/NBP2-33819)



**NBP2-33819**

CXCL8/IL-8 Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS (pH 7.2) and 40% Glycerol

Product Description	
Host	Rabbit
Gene ID	3576
Gene Symbol	CXCL8
Species	Human
Immunogen	This antibody was developed against a recombinant protein corresponding to amino acids: SAKELRCQCIKTYSKPFHPKFIKELRVIESGPHCANTEIIVKLSGDGRELCLDPKEN WVQRVVEKF


Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 0.04-0.4 ug/ml, Immunohistochemistry 1:50 - 1:200, Immunocytochemistry/ Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:50 - 1:200
Application Notes	For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF Fixation Permeabilization: Use PFA/Triton X-100.

**Images**


Western Blot: CXCL8/IL-8 Antibody [NBP2-33819] - Analysis in human cell line EFO-21.




Immunocytochemistry/Immunofluorescence: CXCL8/IL-8 Antibody [NBP2-33819] - Staining of human cell line U-251 MG shows localization to the Golgi apparatus. Antibody staining is shown in green.




Immunohistochemistry-Paraffin: CXCL8/IL-8 Antibody [NBP2-33819] - Staining of human rectum shows moderate positivity in plasma.




Immunohistochemistry-Paraffin: CXCL8/IL-8 Antibody [NBP2-33819] - Staining of human fallopian tube shows no positivity in glandular cells as expected.



Immunohistochemistry-Paraffin: CXCL8/IL-8 Antibody [NBP2-33819] - Staining of human lymph node shows weak to moderate cytoplasmic positivity in a subset of non-germinal center cells.



Immunohistochemistry-Paraffin: CXCL8/IL-8 Antibody [NBP2-33819] - Staining of human placenta shows moderate positivity in plasma.



## Publications

Lee SM, Kim SH, Kim Z et Al. Photodynamic Effects of Topical Photosensitizer, Photodithazine Using Micro-LED for Acne Bacteria Induced Inflammation *Ann Dermatol* 2024-12-01 [PMID: 39623608]

Yu S, Chung S, Ha H et al. Nanotheranostics of Pre Stenotic Vessels By Target Touch On Signaling of Peptide Navigator *Advanced Functional Materials* 2021-12-08 (WB)

Li L, Yao W, Yan S Et al. Pan-Cancer Analysis of Prognostic and Immune Infiltrates for CXCs Cancers 2021-08-18 [PMID: 34439306] (IHC-P, Human)

C Owen-Woods, R Joulia, A Barkaway, L Rolas, B Ma, AF Nottebaum, KP Arkill, M Stein, T Girbl, M Golding, DO Bates, D Vestweber, MB Voisin, S Nourshargh Local microvascular leakage promotes trafficking of activated neutrophils to remote organs *J. Clin. Invest.*, 2020-05-01;0(0):. 2020-05-01 [PMID: 31971917] (IHC, Human)

Fronczek J, Lulf R, Korkmaz Hi, Witte Bi. Analysis of inflammatory cells and mediators in skin wound biopsies to determine wound age in living subjects in forensic medicine *Forensic Science International et al.* 2014-11-22 [PMID: 25506669] (IHC-P, Human)

### Details:

CXCL8/IL-8 antibody used at 1:100 dilution in IHC-P application on human tissues from skin wounds in living subjects in forensic medicine - 4% formaldehyde solution fixation, 4um paraffin sections, endogenous peroxidases blocking with methanol/H<sub>2</sub>O<sub>2</sub> (0.3%) for 30 min, antigen retrieval via boiling of slides for 10 min in Tris-EDTA pH 9, primary incubation for 1 h at room temperature, detection with rabbit Envision - DAB (Fig. 1e and 1f)



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

Error: Subreport could not be shown.

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-33819](http://www.novusbio.com/reviews/submit/NBP2-33819)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

