Product Datasheet

TGF-alpha Antibody (SPM542) NBP2-34371-0.1mg

Unit Size: 0.1 mg

Store at 4C.



Reviews: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34371

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34371



NBP2-34371-0.1mg


TGF-alpha Antibody (SPM542)

0.1 mg	
0.2 mg/ml	
Store at 4C.	
Monoclonal	
SPM542	
0.05% Sodium Azide	
IgM Kappa	
Protein L or PEG purified	
10 mM PBS with 0.05% BSA	
6 kDa	
Product Description	
200ug/ml of antibody purified from Bioreactor Concentrate. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34418) Antibody with azide - store at 2 to 8C.	
Mouse	
7039	
TGFA	
Human, Rabbit, Zebrafish	
This antibody reacts with the TGF alpha and shows no cross-reaction with EGF and the neuropeptide synenkephalin. The staining with this monoclonal antibody is completely blocked by the peptide used for raising this antibody. TGF (aa50) is a growth factor with 33% homology to EGF, binds to EGFR, activates tyrosine phosphorylation of the receptor, and stimulates cell proliferation. It plays a role in tumor initiation by inducing the reversible transformed phenotype.	
A 10-amino acid synthetic peptide (aa34-43; PPVAAAVVSH) from human TGF- alpha (Uniprot: P01135)	
Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin	
Western Blot, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/ml	
Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.	




Images

Western Blot: TGF-alpha Antibody (SPM542) [NBP2-34371] - analysis of lysate from HEK293 cells (30ug) using anti-TGF alpha antibody. Image from verified customer review.



Immunohistochemistry-Paraffin: TGF-alpha Antibody (SPM542) [NBP2-34371] - Formalin-fixed, paraffin-embedded human testicular carcinoma stained with TGF alpha Monoclonal Antibody (SPM542)







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-34371-0.1mg

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-96975-0.5mg	Mouse IgM Kappa Light Chain Isotype Control (MMK)
NBP2-36432-100ug	Human TGF-alpha Recombinant Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34371

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

