Product Datasheet

CD45 Antibody (2B11 + PD7/26) [Alexa Fluor® 350] NBP2-34528AF350

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34528AF350

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34528AF350



NBP2-34528AF350

CD45 Antibody (2B11 + PD7/26) [Alexa Fluor® 350]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClone2B11 + PD7/26Preservative0.05% Sodium AzideIsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionFreeHostMouseGene BD5788Gene SymbolPTPRCSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic unliferential diagnosis of lymphoid tumors from non-hematopoietic unliferential diagnosis of lymphoid tumors from non-hematopoietic undignation on-hematopoietic undifferential diagnosis of lymphoid tumors from non-hematopoietic is useful indignosis of lymphoid tumors from non-hematopoietic is useful indignosis of lymphoid neoplasms. A positive result with this monoclonal antibody is Dressed in regulation of transmembrane signals. Antibody to CD45 is useful indifferential diagnosis of lymphoid tumors from non-hematopoietic undifferential diagnosis of lymphoid tumors from non-hematopoietic indignas is absence does nor trule out a hematopoietic lineage and is present in most being and malignant inon-hematopoietic undifferential diagnosis of lymphoid tumors from Ton-hematopoietic lineage and is present in most being and malignant inon-hematopoietic lineage and is present in most being an dmalignant inon-hematopoietic lineage and is present in most<		[
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Stor at 4C in the dark. Clonality Monoclonal Clone 2B11 + PD7/26 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG1 Kappa Conjugate Alexa Fluor 350 Purity Protein A or G purified Buffer SomM Sodium Borate Product Description Free Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180KDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumor. This antibody is expressed absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic tineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors. <th>Product Information</th> <th></th>	Product Information		
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone 2B11 + PD7/26 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG1 Kappa Conjugate Alexa Fluor 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic uses. The intracellular portions of these molecules have protein phosphates activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differentiated ineoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid moreplasms may lack CD45 (Hodgkin lymphoid regularms or leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors. Immunogen Isolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Unit Size	0.1 ml	
ClonalityMonoclonalClone2B11 + PD7/26Preservative0.05% Sodium AzideIsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180KDa) expressed on hematopoietic cell lines but absent on non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to SD45 is useful in differential diagnosis of lymphoid tumors. The intracellular portions of these molecules have protein phosphatase activity and are involved in negalasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin Jymphoma, sone T-cell lymphoma, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most being and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Concentration		
Clone2B11 + PD7/26Preservative0.05% Sodium AzideIsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180/Da) expressed on hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors. Antibody to Shiphi indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin Lymphomas, and some leukemias) to its absence does not rule out a hematopoietic lineage and is pressent in most berign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Storage	Store at 4C in the dark.	
Preservative0.05% Sodium AzideIsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50MM Sodium BorateProduct DescriptionMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines, normal and malignant non-hematopoietic cell lines, normal and malignant non-hematopoietic lineferentiated in regulation of transmembrane signals. Antioody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic of lymphoid repairs. A positive result with this monoclonal antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most besign and malignant lymphoryphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphoryphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphoryphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphoryphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphoryphosi (2B11); and human peripheral	Clonality	Monoclonal	
IsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180KDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic cull are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Clone	2B11 + PD7/26	
ConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is nighly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Preservative	0.05% Sodium Azide	
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphoma, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Isotype	IgG1 Kappa/IgG1 Kappa	
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic issues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Conjugate	Alexa Fluor 350	
Product Description Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors. Immunogen Isolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Purity	Protein A or G purified	
HostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Buffer	50mM Sodium Borate	
Gene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of Jymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Product Description		
Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphoma, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Host	Mouse	
SpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Gene ID	5788	
MarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Gene Symbol	PTPRC	
Specificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Species	Human	
 comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors. Isolated neoplastic cells from T cell lymphoma (2B11); and human peripheral 	Marker	Leucocyte Marker	
	Specificity/Sensitivity	comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most	
	Immunogen		



N	otes	
Ν	otes	

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details

Applications	Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-ready
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Images

CD45 Antibody (2B11 + PD7/26) [Alexa Fluor® 350] [NBP2-34528AF350] - Vial of Alexa Fluor 350 conjugated antibody. Alexa Fluor 350 is optimally excited at 346 nm by the UV laser (350 or 355 nm) and has an emission maximum of 442 nm.

The state of the s			
in and		@ २ ८०	
	Alexa Fluc	or® 350	
Safety Marca	LASER (nm)	FILTER	
	UV (350)	450/45	
	EXCITATION MAX (nm)	EMISSION MAX (nm)	
	346	442	
A NOVUS			
CAUTION -Recearch Use Only			





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-34528AF350

NBP2-51591-0.02mg	Recombinant Human CD45 His Protein
210-TA-005	TNF-alpha [Unconjugated]
1430-CD-050	CD45 [Unconjugated]
M6000B-1	IL-6 [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34528AF350

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

