## **Product Datasheet**

### CD45 Antibody (2B11 + PD7/26) [Alexa Fluor® 700] NBP2-34528AF700

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34528AF700

Updated 10/23/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34528AF700



#### NBP2-34528AF700

CD45 Antibody (2B11 + PD7/26) [Alexa Fluor® 700]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClone2B11 + PD7/26Preservative0.05% Sodium AzideIsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionFreesonal ConjugateHostMouseGene ID5788Gene SymbolPTPRCSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoletic undifferential diagnosis of lymphoid tumors from non-hematopoletic undifferential diagnosis of lymphoid tumors from non-hematopoletic undifferential diagnosis of lymphoid tumors from non-hematopoletic undifferential diagnosis of lymphoid tumors. The intracellular portions of species have protein phosphates activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful and diagnosis of lymphoid tumors from non-hematopoletic undifferential diagnosis of lymphoid tumors. The intracellular portions of lymphoid tumors from non-hematopoletic undifferential diagnosis of lymphoid tumors. The sintacellular portions of lymphoid tumors from non-hematopoletic undifferential diagnosis of lymphoid tumors. The sintacellular portions of lymphoid tumors from non-hematopoletic undifferential diagnosis of lymphoid tumors. The sintacellular portions absence does not rule out a hematolymphoid tumor. This antibody is presessed in most being an antignal tymphores as well as	<b>, , ,</b>	
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClone2B11 + PD7/26Preservative0.05% Sodium AzideIsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description5788Gene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180KDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell inses, normal and malignant non-hematopoietic undifferential diagois of lymphoid more from soft from soft mono-hematopoietic cell lines, normal and malignant non-hematopoietic undifferential diagois of lymphoid romora. A positive result with this monoclonal are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagois of lymphoid utomora. This antibody is expressed almost exclusively by cells of hematopoietic tundifferentiated meoplasms. A positive result with this monoclonal and insegue and is present in most beingn and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalClone2B11 + PD7/26Preservative0.05% Sodium AzideIsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description5788Gene ID5788Gene SymbolPTPRCSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180/ka) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphnates activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differentiated neoplastic cells from T-cell lymphoma (2B11); and human peripheralImmunogenIsolated neoplastic cells from T-cell lymphoma (2B11); and human peripheral	Unit Size	0.1 ml
ClonalityMonoclonalClone2B11 + PD7/26Preservative0.05% Sodium AzideIsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines, normal and malignant non-hematopoietic issues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid origin. Certain types of lymphoid neoplasms. A positive result with this monoclonal antibody is expressed almost exclusively by cells of hematopoietic lineage and is pressent in most bergin and malignant tony. The sand some leukemiae) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is pressent in most bergin and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Concentration	
Clone2B11 + PD7/26Preservative0.05% Sodium AzideIsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionKappaHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines, normal and malignant non-hematopoietic cell lines, normal and malignant non-hematopoietic unifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid utmors. An positive result with this monoclonal antibody is expressed antost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic or lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic fungage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Clonality	Monoclonal
IsotypeIgG1 Kappa/IgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180KDa) expressed on hematopoietic cisues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasm say lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is pressent in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Clone	2B11 + PD7/26
ConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most bering and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Isotype	IgG1 Kappa/IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Conjugate	Alexa Fluor 700
Product DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180KDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopynetic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Purity	Protein A or G purified
HostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Buffer	50mM Sodium Borate
Gene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of Jymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of Iymphoid or myeloid origin. Certain types of Iymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Product Description	
Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Host	Mouse
SpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Gene ID	5788
MarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Gene Symbol	PTPRC
Specificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral	Species	Human
<ul> <li>comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.</li> <li>Immunogen</li> </ul>	Marker	Leucocyte Marker
	Specificity/Sensitivity	comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most
	Immunogen	

www.novusbio.com



EXCITATION MAX (nm) EMISSION MAX (nm)

720

696

	1 age 2 01 3 4.20.1 Optialed 10/23/2024
Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-ready
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-ready
Images	
	ior 700 conjugated antibody. Alexa Fluor mm by the Red laser (633 or 640 nm) and





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-34528AF700

NBP2-51591-0.02mg	Recombinant Human CD45 His Protein
210-TA-005	TNF-alpha [Unconjugated]
1430-CD-050	CD45 [Unconjugated]
M6000B-1	IL-6 [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34528AF700

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

