Product Datasheet CD45 Antibody (2B11 + PD7/26) [HRP] NBP2-34528H

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34528H

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34528H



NBP2-34528H

CD45 Antibody (2B11 + PD7/26) [HRP]

Product Information Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clonality Monoclonal Clonality Monoclonal Clone 2B11 + PD7/26 Preservative No Preservative Isotype IgG1 Kappa/IgG1 Kappa Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 1180k/20) expressed on hematopoietic cell lines is usabed on inordopoietic undifferential diagnosis of lymphonid neoplasms may lack CD45 Hodgkin lymphoma, some T-cell lymphom		
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone 2B11 + PD7/26 Preservative No Preservative tsotype IgG1 Kappa/IgG1 Kappa Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 5788 Gene Symbol PTPRC Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180KDa) expressed on hematopoletic cell lines but absent on non-hematopoletic cell lines, normal and malignant non-hematopoletic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmerbrane signals. Antibody to CD45 is leveful in differentiated necoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid tumors from non-hematopoletic usues. The instantbody is expressed almost exclusively by cells of thematopoletic lineage and is present in most being a	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalClone2B11 + PD7/26PreservativeNo PreservativeIsotypeIgG1 Kappa/IgG1 KappaConjugateHRPPurityProtein A or G purifiedButferPBSProduct DescriptionDescriptionDescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID5788Gene SymbolPTPRCSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (S20, 205, 190, 180Kba) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic undifferential diagnosis of lymphoid tumors from non-hematopoietic undifferential diagnosis of lymphoid tumors from non-hematopoietic molyced in regulation of transmembrane signals. Antibody is highly indicative of lymphoid tumors, sont or-cell lymphoma, sont or-cell lymphore is loss of sabence does not rule out a hematolympic line and sont persent in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (SDT); and human peripheral bloid or myeloid origin. Certain types of lymphoid tumors is absence does not rule out a hematolympic line and sont persent in most exbensively by cells of hematopoietic lineage and is present in most exbensively by cells of hematopoietic l	Unit Size	0.1 ml
Clonality Monoclonal Clone 2B11 + PD7/26 Preservative No Preservative Isotype IgG1 Kappa/IgG1 Kappa Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180, 180, 200) expressed on hematopoietic cull ines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic cull issues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differentiated ineoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid tumors from non-hematopoietic culliferentiated and aposis of lymphoid tumors. The instracellular portions of these hotheautophympic tumor. This	Concentration	
Clone 2B11 + PD7/26 Preservative No Preservative Isotype IgG1 Kappa/IgG1 Kappa Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 5783 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180 KDa) expressed on hematopoietic cell lines, normal and malignant non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is buffy indicative of lymphoid or pyneloid origin. Certain types of lymphoid curgolasmes. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is userful and malignast so: Lassen does not rule out a hematopoietic lineage and is present in most being and malignant sort. Neeplasms may lack CD45 (Hodgkin lymphoid neoplasms curgolises the CD45 (Hodgkin lymphoid set pressed and neoplasms and y lack CD45 (Hodgkin lymphore, so of rule out a hematopoietic lineage and is present in most bering and malignant lymphores of ly	Storage	Store at 4C in the dark.
Preservative No Preservative Isotype IgG1 Kappa/IgG1 Kappa Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in negulation of transmerbrane signals. Antibody to CD45 is useful in differentiated negolasms. A positive result with this monoclonal antibody is highly indicative of lymphoid englasm servers and some leukemias) so its absence does not rule out a hematophynehold tumor. This antibody is spressed atmost exclusively by cells of hematopoietic tissues and some leukemias) so its absence does not rule out a hematophynehold tumor. This antibody is spressed atmost exclusively by cells of hematopoietic tissues. The intracellular portion and malignant lymphorynes and some leukemias) so its absence does not rule out a hematop	Clonality	Monoclonal
IsotypeIgG1 Kappa/IgG1 KappaConjugateHRPPurityProtein A or G purifiedBufferPBSProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phold tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid some leukemias) so its absence does not rule out a hematopoietic timage and is present in most bening and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26)Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry. Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry.	Clone	2B11 + PD7/26
Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoletic tissues. The intracellular portions of flymphoid tumors from non-hematopoletic undifferentiated diagnosis of lymphoid tumors from non-hematopoletic undifferentiated diagnosis of lymphoid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoletic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors. Immunogen Isolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready <th>Preservative</th> <th>No Preservative</th>	Preservative	No Preservative
Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180KDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differentiated neoplasms. A positive result with this monoclonal antibody is bighly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphoma, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors. Immunogen Isolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26) Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry- Paraffin, CyTOF-ready Recomme	Isotype	IgG1 Kappa/IgG1 Kappa
Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic issues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody is CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is supressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant torypes of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant tymphocytes as well as plasma cell precursors. Immunogen Isolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26) Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry, I	Conjugate	HRP
Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell ines normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic used in regulation of transmembrane signals. Antibody to SL45 is useful in differential diagnosis of lymphoid tumors. This antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors. Immunogen Isolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26) Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Pararaffin, CyTOF-read	Purity	Protein A or G purified
DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic lissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26)Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry. Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry.	Buffer	PBS
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26)Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry	Product Description	
Gene ID5788Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukernias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26)Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry. Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-ready	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene SymbolPTPRCSpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26)Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry	Host	Mouse
SpeciesHumanMarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26)Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry	Gene ID	5788
MarkerLeucocyte MarkerSpecificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26)Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry	Gene Symbol	PTPRC
Specificity/SensitivityRecognizes the CD45 leukocyte common antigen (LCA) family which is comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26)Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry,	Species	Human
comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most benign and malignant lymphocytes as well as plasma cell precursors.ImmunogenIsolated neoplastic cells from T cell lymphoma (2B11); and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26)Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry	Marker	Leucocyte Marker
blood lymphocytes maintained in T cell growth factor (PD7/26) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-	Specificity/Sensitivity	comprised of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portions of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms. A positive result with this monoclonal antibody is highly indicative of lymphoid or myeloid origin. Certain types of lymphoid neoplasms may lack CD45 (Hodgkin lymphoma, some T-cell lymphomas, and some leukemias) so its absence does not rule out a hematolymphoid tumor. This antibody is expressed almost exclusively by cells of hematopoietic lineage and is present in most
ApplicationsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-	Immunogen	
Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-	Product Application Details	
	Applications	
	Recommended Dilutions	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-34528H

NBP2-51591-0.02mg	Recombinant Human CD45 His Protein
210-TA-005	TNF-alpha [Unconjugated]
1430-CD-050	CD45 [Unconjugated]
M6000B-1	IL-6 [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34528H

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

