Product Datasheet

EpCAM/TROP1 Antibody (323/A3) [mFluor Violet 500 SE] NBP2-34532MFV500

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34532MFV500

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34532MFV500



NBP2-34532MFV500

EpCAM/TROP1 Antibody (323/A3) [mFluor Violet 500 SE]

Product	Information	
Toduot	mormation	

epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalClone323/A3Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugatemFluor Violet 500 SEPurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID4072Gene SymbolEPCAMSpeciesHuman, Rat (Negative)Reactivity NotesDoes not react with Rat, Feline or Canine.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular achesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and ehoaccinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody has been used to distinguish a adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody has been used to distinguish a adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody has been used to distinguish a adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody has b	Unit Size	0.1 ml
ClonalityMonoclonalClone323/A3Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugatemFluor Violet 500 SEPurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4072Gene SymbolEPCAMSpeciesHuman, Rat (Negative)Reactivity NotesDoes not react with Rat, Feline or Canine.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43XDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia adhoroarcinoman from pleural mesothelloma and hepatcellular carcinoma. This antibody has been used to distinguish adhoroarcinoma from pleural mesothelloma and hepatcellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelloma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready	Concentration	•
Clone323/A3Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugatemFluor Violet 500 SEPurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description4072HostMouseGene ID4072Gene SymbolEPCAMSpeciesHuman, Rat (Negative)Reactivity NotesDoes not react with Rat, Feline or Canine.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific anigen (ESA), or epithelial cellular adhesion moleculle (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugatemFluor Violet 500 SEPurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID4072Gene SymbolEPCAMSpeciesHuman, Rat (Negative)Reactivity NotesDoes not react with Rat, Feline or Canine.MarkerEgGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on 	Clonality	Monoclonal
IsotypeIgG1 KappaConjugatemFluor Violet 500 SEPurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID4072Gene SymbolEPCAMSpeciesHuman, Rat (Negative)Reactivity NotesDoes not react with Rat, Feline or Canine.MarkerEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cellustrace in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry. Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry.Paraffin, CyTOF-ready	Clone	323/A3
ConjugatemFluor Violet 500 SEPurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID4072Gene SymbolEPCAMSpeciesHuman, Rat (Negative)Reactivity NotesDoes not react with Rat, Feline or Canine.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM) is astocarcinoma. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on droud recovery may vary from the stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry/Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Paraffin, CyTOF-ready	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description 4072 Host Mouse Gene ID 4072 Gene Symbol EPCAM Species Human, Rat (Negative) Reactivity Notes Does not react with Rat, Feline or Canine. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and evast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Immunogen MCF-7 human breast cancer cells Notes mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry, Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistoc	Isotype	IgG1 Kappa
Buffer 50mM Sodium Borate Product Description Mouse Host Mouse Gene ID 4072 Gene Symbol EPCAM Species Human, Rat (Negative) Reactivity Notes Does not react with Rat, Feline or Canine. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and evast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Immunogen MCF-7 human breast cancer cells Notes mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry, Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immu	Conjugate	mFluor Violet 500 SE
Product Description Mouse Host Mouse Gene ID 4072 Gene Symbol EPCAM Species Human, Rat (Negative) Reactivity Notes Does not react with Rat, Feline or Canine. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and avat majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Immunogen MCF-7 human breast cancer cells Notes mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Paraffin, CyTOF-ready	Purity	Protein A or G purified
HostMouseGene ID4072Gene SymbolEPCAMSpeciesHuman, Rat (Negative)Reactivity NotesDoes not react with Rat, Feline or Canine.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry, Immunohistochemistry, Immunohistochemistry, CyTOF-ready	Buffer	50mM Sodium Borate
Gene ID4072Gene SymbolEPCAMSpeciesHuman, Rat (Negative)Reactivity NotesDoes not react with Rat, Feline or Canine.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry-Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry. Paraffin, CyTOF-ready	Product Description	
Gene SymbolEPCAMSpeciesHuman, Rat (Negative)Reactivity NotesDoes not react with Rat, Feline or Canine.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry. Immunofluorescence, Immunohistochemistry/Immunofluorescence, Immunohistochemistry. Paraffin, CyTOF-ready	Host	Mouse
Species Human, Rat (Negative) Reactivity Notes Does not react with Rat, Feline or Canine. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Immunogen MCF-7 human breast cancer cells Notes mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Paraffin, CyTOF-ready	Gene ID	4072
Reactivity NotesDoes not react with Rat, Feline or Canine.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia ad a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, 	Gene Symbol	EPCAM
MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry. Immunohistochemistry, Immunocytochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry/Immunofluorescence, Immunohistochemistry. Paraffin, CyTOF-ready	Species	Human, Rat (Negative)
Specificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, CyTOF-ready	Reactivity Notes	Does not react with Rat, Feline or Canine.
epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenMCF-7 human breast cancer cellsNotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry/Immunofluorescence, Immunohistochemistry/Immunofluorescence, Immunohistochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistr	Marker	Epithelial Marker
NotesmFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry/Immunofluorescence, Immunohistochemistry, Paraffin, CyTOF-ready	Specificity/Sensitivity	CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from
demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.Product Application DetailsApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunohistochemistry, Immunohistochemistry/Immunohistochemistry, Immunocytochemistry/Immunohistochemistry, Immunohistochemistry/Immunohistochemistry, Immunohistochemistry/Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready	Immunogen	MCF-7 human breast cancer cells
ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-readyRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready	Notes	demand. Actual recovery may vary from the stated volume of this product. The
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready	Product Application Details	
Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready	Applications	
Application Notes Optimal dilution of this antibody should be experimentally determined.	Recommended Dilutions	Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin,
	Application Notes	Optimal dilution of this antibody should be experimentally determined.



Images

EpCAM/TROP1 Antibody (323/A3) [mFluor Violet 500 SE] - Vial of mFluor Violet 500 conjugated antibody. mFluor Violet 500 is optimally excited at 410 nm by the Violet laser (405 nm) and has an emission maximum of 501 nm.



www.novusbio.com





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 novus@novusbio.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: technical@novusbio.com Orders: orders@novusbio.com General: novus@novusbio.com

Products Related to NBP2-34532MFV500

NBP2-52190-0.05mg	Recombinant Human EpCAM/TROP1 His Protein
202-IL-010	IL-2 [Unconjugated]
960-EP-050	EpCAM/TROP1 [Unconjugated]
MEP00B	Erythropoietin/EPO [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34532MFV500

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

