Product Datasheet

Golgi Complex Antibody (371-4) [Janelia Fluor® 646] NBP2-34534JF646

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34534JF646

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34534JF646



NBP2-34534JF646

Unit Size Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone 371-4 Preservative 10.05% Sodium Azide 1sotype 1gG1 Kappa Conjugate Janelia Fluor 646 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker or Human Cells Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flatened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of taltened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of taltened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though the pending on where they are destined to reside, Several spherical vesicles that have budded off of the Golgi are present surro	Golgi Complex Antibody (371-4) [Janelia Fluor® 646]		
Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark.	Product Information		
Storage Store at 4C in the dark. Clonality Monoclonal Clone 371-4 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Janelia Fluor 646 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker of Human Cells Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant eclls. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells in xenographic model research. It reacts specifically with human cells in xenographic model research. It reacts specifically with human cells in the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sack shrow as disternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. Immunogen SU-DHL-1 large cell lymphoma cells. Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Unit Size	0.1 ml	
Clonality Monoclonal Clone 371-4 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Janelia Fluor 646 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellual rractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flatened, membrane bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. Immunogen SU-DHL-1 large cell lymphoma cells. Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Concentration	· ·	
Clone 371-4 Preservative 0.05% Sodium Azide Isotype 1gG1 Kappa Janelia Fluor 646 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Marker for Human Cells Specificity/Sensitivity Disconsistivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in human cells only. It can be used to stain the Golgi complex in human cells only. It can be used to stain the Golgi complex in human cells only. It can be used to stain the Golgi complex in human cells only. It can be used to stain the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of usions enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. Immunogen SU-DHL-1 large cell lymphoma cells. Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Storage	Store at 4C in the dark.	
Preservative 0.05% Sodium Azide Isotype	Clonality	Monoclonal	
Isotype IgG1 Kappa Conjugate Janelia Fluor 646 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cisface, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. Immunogen SU-DHL-1 large cell lymphoma cells. Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Clone	371-4	
Conjugate Janelia Fluor 646 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. Immunogen SU-DHL-1 large cell lymphoma cells. Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Preservative	0.05% Sodium Azide	
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sack known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. Immunogen SU-DHL-1 large cell lymphoma cells. Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Isotype	IgG1 Kappa	
Buffer SomM Sodium Borate	Conjugate	Janelia Fluor 646	
Product Description	Purity	Protein A or G purified	
Host	Buffer	50mM Sodium Borate	
Reactivity Notes Does not react with Mouse or Rat.	Product Description		
Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells This monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. Immunogen SU-DHL-1 large cell lymphoma cells. Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Host	Mouse	
MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.ImmunogenSU-DHL-1 large cell lymphoma cells.NotesSold under license from the Howard Hughes Medical Institute, Janelia Research Campus.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Species	Human, Mouse (Negative), Rat (Negative)	
Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. Immunogen SU-DHL-1 large cell lymphoma cells. Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Reactivity Notes	Does not react with Mouse or Rat.	
complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. Immunogen SU-DHL-1 large cell lymphoma cells. Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Marker	Marker for Human Cells	
Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Specificity/Sensitivity	complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	
Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Immunogen	SU-DHL-1 large cell lymphoma cells.	
Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Notes	· ·	
Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	Product Application Details		
Immunohistochemistry-Paraffin	Applications		
Application Notes Optimal dilution of this antibody should be experimentally determined.	Recommended Dilutions		
	Application Notes	Optimal dilution of this antibody should be experimentally determined.	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34534JF646

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

