Product Datasheet

CD5 Antibody (C5/473 + CD5/54/F6) [Alexa Fluor® 532] NBP2-34583AF532

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34583AF532

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34583AF532



NBP2-34583AF532

CD5 Antibody (C5/473 + CD5/54/F6) [Alexa Fluor® 532]

Product Information Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone C5/473 + CD5/54/F6 Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas. Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplastic B-
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone C5/473 + CD5/54/F6 Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone C5/473 + CD5/54/F6 Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Clonality Monoclonal Clone C5/473 + CD5/54/F6 Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Clone C5/473 + CD5/54/F6 Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Isotype IgG1 Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Conjugate Alexa Fluor 532 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Product Description Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Gene Symbol CD5 Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Gene Symbol Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Species Human, Rat, Bovine, Rabbit Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas.
cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SL mantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymphoma. CD5 aberrant expression is useful in making a diagnosis of mature T-cell neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of othe B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also very useful in differentiating among mature small lymphoid cell malignancies. In addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) from thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.
Immunogen Human CD5 recombinant protein (C5/473); A synthetic peptide from the intracellular region of human CD5 (CD5/54/F6)



Notes

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details	
Applications	ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34583AF532

NBP1-97005AF532 Mouse IgG1 Isotype Control (MG1) [Alexa Fluor® 532]

NBP2-38900PEP CD5 Recombinant Protein Antigen

6507-IL-010/CF IL-4 [Unconjugated]

1636-CD-050 CD5

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34583AF532

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

