Product Datasheet

CEACAM5/CD66e Antibody (C66/261) [Janelia Fluor® 585] NBP2-34586JF585

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34586JF585

Updated 12/12/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34586JF585



NBP2-34586JF585

CEACAM5/CD66e Antibody (C66/261) [Janelia Fluor® 585]

_
ee the vial label for concentration. If unlisted please contact technical
C in the dark.
al
dium Azide
ра
uor 585
or G purified
dium Borate
5
rody recognizes proteins of 80-200kDa, identified as different members mily. CEA is synthesized during development in the fetal gut and is red in increased amounts in intestinal carcinomas and several other his monoclonal antibody does not react with nonspecific cross-reacting NCA) and with human polymorphonuclear leucocytes. It shows no with a variety of normal tissues and is suitable for staining of paraffin tissues. CEA is not found in benign glands, stroma, or malignant cells. Antibody to CEA is useful in detecting early foci of gastric a and in distinguishing pulmonary adenocarcinomas (60-70% are CEA eural mesotheliomas (rarely or weakly CEA+). Anti-CEA positivity is denocarcinomas from the lung, colon, stomach, esophagus, pancreas, er, urachus, salivary gland, ovary, and endocervix.
rody recognizes proteins of 80-200kDa, identified as different members mily. CEA is synthesized during development in the fetal gut and is red in increased amounts in intestinal carcinomas and several other his monoclonal antibody does not react with nonspecific cross-reacting NCA) and with human polymorphonuclear leucocytes. It shows no with a variety of normal tissues and is suitable for staining of paraffin tissues. CEA is not found in benign glands, stroma, or malignant cells. Antibody to CEA is useful in detecting early foci of gastric a and in distinguishing pulmonary adenocarcinomas (60-70% are CEA eural mesotheliomas (rarely or weakly CEA+). Anti-CEA positivity is denocarcinomas from the lung, colon, stomach, esophagus, pancreas,
rody recognizes proteins of 80-200kDa, identified as different members mily. CEA is synthesized during development in the fetal gut and is red in increased amounts in intestinal carcinomas and several other his monoclonal antibody does not react with nonspecific cross-reacting NCA) and with human polymorphonuclear leucocytes. It shows no with a variety of normal tissues and is suitable for staining of paraffin tissues. CEA is not found in benign glands, stroma, or malignant cells. Antibody to CEA is useful in detecting early foci of gastric a and in distinguishing pulmonary adenocarcinomas (60-70% are CEA eural mesotheliomas (rarely or weakly CEA+). Anti-CEA positivity is denocarcinomas from the lung, colon, stomach, esophagus, pancreas, er, urachus, salivary gland, ovary, and endocervix.
rody recognizes proteins of 80-200kDa, identified as different members mily. CEA is synthesized during development in the fetal gut and is red in increased amounts in intestinal carcinomas and several other his monoclonal antibody does not react with nonspecific cross-reacting NCA) and with human polymorphonuclear leucocytes. It shows no with a variety of normal tissues and is suitable for staining of paraffin tissues. CEA is not found in benign glands, stroma, or malignant cells. Antibody to CEA is useful in detecting early foci of gastric a and in distinguishing pulmonary adenocarcinomas (60-70% are CEA eural mesotheliomas (rarely or weakly CEA+). Anti-CEA positivity is denocarcinomas from the lung, colon, stomach, esophagus, pancreas, er, urachus, salivary gland, ovary, and endocervix.
rody recognizes proteins of 80-200kDa, identified as different members mily. CEA is synthesized during development in the fetal gut and is red in increased amounts in intestinal carcinomas and several other his monoclonal antibody does not react with nonspecific cross-reacting NCA) and with human polymorphonuclear leucocytes. It shows no with a variety of normal tissues and is suitable for staining of paraffin tissues. CEA is not found in benign glands, stroma, or malignant cells. Antibody to CEA is useful in detecting early foci of gastric a and in distinguishing pulmonary adenocarcinomas (60-70% are CEA eural mesotheliomas (rarely or weakly CEA+). Anti-CEA positivity is denocarcinomas from the lung, colon, stomach, esophagus, pancreas, er, urachus, salivary gland, ovary, and endocervix.
rody recognizes proteins of 80-200kDa, identified as different members mily. CEA is synthesized during development in the fetal gut and is red in increased amounts in intestinal carcinomas and several other his monoclonal antibody does not react with nonspecific cross-reacting NCA) and with human polymorphonuclear leucocytes. It shows no with a variety of normal tissues and is suitable for staining of paraffin tissues. CEA is not found in benign glands, stroma, or malignant cells. Antibody to CEA is useful in detecting early foci of gastric a and in distinguishing pulmonary adenocarcinomas (60-70% are CEA eural mesotheliomas (rarely or weakly CEA+). Anti-CEA positivity is denocarcinomas from the lung, colon, stomach, esophagus, pancreas, er, urachus, salivary gland, ovary, and endocervix. The paraffin tissues. CEA is not found in benign glands, stroma, or malignant cells. Antibody to CEA is useful in detecting early foci of gastric a and in distinguishing pulmonary adenocarcinomas (60-70% are CEA eural mesotheliomas (rarely or weakly CEA+). Anti-CEA positivity is denocarcinomas from the lung, colon, stomach, esophagus, pancreas, er, urachus, salivary gland, ovary, and endocervix. The paraffin tissues.
1





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34586JF585

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

