# **Product Datasheet**

# CD8 Antibody (C8/468) [Allophycocyanin] NBP2-34589APC

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34589APC

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34589APC



#### NBP2-34589APC

| CD8 Antibody (C8/468) [Allophycocyanin]   |  |
|---|--|
|   |  |
| 0.1 ml  |  |
| Please see the vial label for concentration. If unlisted please contact technical services.   |  |
| Store at 4C in the dark.  |  |
| Monoclonal  |  |
| C8/468  |  |
| 0.05% Sodium Azide  |  |
| IgG1 Kappa  |  |
| Allophycocyanin   |  |
| Protein A or G purified   |  |
| PBS   |  |
|   |  |
| This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.   |  |
| Mouse   |  |
| 925   |  |
| CD8A  |  |
| Human   |  |
| Cytotoxic- & Suppressor T-Cell Marker   |  |
| CD8 Antibody (C8/468) [Allophycocyanin] recognizes CD8, a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha/beta) or as a homodimer (CD8 alpha/alpha). A majority of thymocytes and a subpopulation of mature T cells and NK cells express CD8a. CD8 binds to MHC class 1 and through its association with protein tyrosine kinase p56lck plays a role in T cell development and activation of mature T cells. For mature T-cells, CD4 and CD8 are mutually exclusive, so anti-CD8 is generally used in conjunction with anti-CD4. It is a useful marker for distinguishing helper/inducer T-lymphocytes, and most peripheral T-cell lymphomas are CD4+/CD8 Anaplastic large cell lymphoma is usually CD4+ and CD8-, and in T-lymphoblastic lymphoma/leukemia, CD4 and CD8 are often co-expressed. CD8 is also found in littoral cell angioma of the spleen. |  |
| Human CD8 recombinant protein (Uniprot: P01732)   |  |
|   |  |
| Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready   |  |
| Flow Cytometry, Immunohistochemistry, Immunocytochemistry/<br>Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready  |  |
|   |  |



# **Images**

CD8 Antibody (C8/468) [Allophycocyanin] [NBP2-34589APC] - Vial of APC conjugated antibody. APC is optimally excited at 650 nm by the Red laser (633 or 640 nm) and has an emission maximum of 660 nm.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NBP2-34589APC**

IC002A Mouse IgG1 Isotype Control (11711) [Allophycocyanin]

NBP2-36743PEP CD8 Recombinant Protein Antigen

210-TA-005 TNF-alpha [Unconjugated]

9358-CD-050 CD8 [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34589APC

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

