# **Product Datasheet**

# CD5 Antibody (CD5/54/F6) [DyLight 650] NBP2-34593C

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34593C

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34593C



# NBP2-34593C

CD5 Antibody (CD5/54/F6) [DyLight 650]

Product Information	
Concentration  Please see the vial label for concentration. If unlisted please contact tec services.  Storage  Store at 4C in the dark.  Clonality  Monoclonal  Clone  CD5/54/F6  Preservative  0.05% Sodium Azide  Isotype  IgC1 Kappa  Conjugate  DyLight 650  Purity  Protein A or G purified  Buffer  SomM Sodium Borate  Product Description  This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the ustated on the datasheet.  Host  Mouse  Gene ID  921  Gene Symbol  CD5  Species  Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, R. Marker  Mantle Cell Lymphoma Marker  Specificity/Sensitivity  Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell and Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplas cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CD5 abertant expression is useful in making a diagnosis of mature T-ce neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel B-cell marker, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies. addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) from thymoma (-). Anti-CD5 does not react with granulocytes or monocytes. Immunogen  Immunogen	
Storage Store at 4C in the dark.  Clonality Monoclonal  Clone CD5/54/F6  Preservative 0.05% Sodium Azide  Isotype IgG1 Kappa  Conjugate DyLight 650  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Description This conjugate is made on demand. Actual recovery may vary from the avolume of this product. The volume will be greater than or equal to the datasheet.  Host Mouse  Gene ID 921  Gene Symbol CD5  Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, R. Marker Mantle Cell Lymphoma Marker  Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell at Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplas cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (Cmantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymph CD5 aberrant expression is useful in making a diagnosis of mature T-ceneoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell marker, especially one that includes ancibe 23s. Anti-CD5 is abe useful in differentiating among mature small lymphoid cell malignancies, addition, anti-CD5 does not react with granulocytes or monocytes. Immunogen  Recombinant fragment (around aa450-495) of human CD5 protein (inter	
Clonality Monoclonal Clone CD5/54/F6 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Description This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the ustated on the datasheet.  Host Mouse Gene ID 921 Gene Symbol CD5 Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, R. Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell at Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplas cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (and a subset (~10%) of diffuse large B-cell lympl CD5 aberrant expression is useful in making a diagnosis of mature T-ce neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel de B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) frithymoma (-). Anti-CD5 can be used in distinguishing thymic carcinoma (+) frithymoma (-). Anti-CD5 can be used to react with granulocytes or monocytes.	nical
Clone CD5/54/F6 Preservative Jeg1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer SomM Sodium Borate  Product Description This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the ustated on the datasheet.  Host Mouse Gene ID Gene Symbol CD5 Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, R. Marker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell at Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplas cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CD5 aberrant expression is useful in making a diagnosis of mature T-ce neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD52. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies addition, anti-CD5 does not react with granulocytes or monocytes.  Immunogen Recombinant fragment (around aa450-495) of human CD5 protein (intra	
Preservative  Isotype  IgG1 Kappa  Conjugate  DyLight 650  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the ustated on the datasheet.  Host  Mouse  Gene ID  921  Gene Symbol  CD5  Species  Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, R. Marker  Mantle Cell Lymphoma Marker  Specificity/Sensitivity  Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell at Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplass cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (Cmantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymph CD5 aberrant expression is useful in making a diagnosis of mature T-ce neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies. addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) from thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen  Recombinant fragment (around aa450-495) of human CD5 protein (intra	
Isotype	
Conjugate Purity Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Description  This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the ustated on the datasheet.  Mouse  Gene ID  921  Gene Symbol  CD5  Species  Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, R. Marker  Mantle Cell Lymphoma Marker  Specificity/Sensitivity  Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell an Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplas cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (C mantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymph CD5 aberrant expression is useful in making a diagnostic in CLL/SLL within a panel B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies. addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) frumoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen  Recombinant fragment (around aa450-495) of human CD5 protein (intra	
Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the ustated on the datasheet.  Host Mouse  Gene ID 921  Gene Symbol CD5  Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, R. Marker Mantle Cell Lymphoma Marker  Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell and Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplas cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (C mantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymph CD5 aberrant expression is useful in making a diagnosis of mature T-ce neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies. addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) fir thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen Recombinant fragment (around aa450-495) of human CD5 protein (intra	
Product Description  Description  This conjugate is made on demand. Actual recovery may vary from the experiment of this product. The volume will be greater than or equal to the distance on the datasheet.  Host  Mouse  Gene ID  Gene Symbol  CD5  Species  Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, Ramarker  Mantle Cell Lymphoma Marker  Specificity/Sensitivity  Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell at Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplas cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CD5 aberrant expression is useful in making a diagnosis of mature T-ce neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) fir thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen  Recombinant fragment (around aa450-495) of human CD5 protein (intra	
Product Description  This conjugate is made on demand. Actual recovery may vary from the explain volume of this product. The volume will be greater than or equal to the ustated on the datasheet.  Mouse  Gene ID  921  Gene Symbol  CD5  Species  Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, Rather  Mantle Cell Lymphoma Marker  Specificity/Sensitivity  Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloodlymphocytes. In lymph nodes, the main reactivity is observed in T cell and Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplast cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CD5 aberrant expression is useful in making a diagnosis of mature T-cent neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies.  B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies. addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) from thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen  Recombinant fragment (around aa450-495) of human CD5 protein (intra	
This conjugate is made on demand. Actual recovery may vary from the evolume of this product. The volume will be greater than or equal to the distance on the datasheet.  Mouse  Gene ID  921  Gene Symbol  CD5  Species  Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, Ranker  Mantle Cell Lymphoma Marker  Specificity/Sensitivity  Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell ar Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplas cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (C mantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymph CD5 aberrant expression is useful in making a diagnosis of mature T-ce neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD25 is also useful in differentiating among mature small lymphoid cell malignancies, addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) frummunogen  Recombinant fragment (around aa450-495) of human CD5 protein (intra	
volume of this product. The volume will be greater than or equal to the ustated on the datasheet.  Mouse  Gene ID  921  Gene Symbol  CD5  Species  Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, Ramarker  Mantle Cell Lymphoma Marker  Specificity/Sensitivity  Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell and Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplas cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CD5 aberrant expression is useful in making a diagnosis of mature T-ceneoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD5 is addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) from thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen  Recombinant fragment (around aa450-495) of human CD5 protein (intra	
Gene Symbol CD5 Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, Ramarker Mantle Cell Lymphoma Marker Specificity/Sensitivity Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell are Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplass cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphomantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymph CD5 aberrant expression is useful in making a diagnosis of mature T-ceneoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) for thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen  Recombinant fragment (around aa450-495) of human CD5 protein (intra	
Gene Symbol  CD5  Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, Rather  Mantle Cell Lymphoma Marker  Specificity/Sensitivity  Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell ar Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplass cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (Cmantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymph CD5 aberrant expression is useful in making a diagnosis of mature T-ceneoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) from thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen  Recombinant fragment (around aa450-495) of human CD5 protein (intra	
Species  Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Monkey, Rather  Mantle Cell Lymphoma Marker  Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell and Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplass cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CD5 aberrant expression is useful in making a diagnosis of mature T-center neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies. addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) from thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen	
Marker  Mantle Cell Lymphoma Marker  Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell at Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplas cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (C mantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymph CD5 aberrant expression is useful in making a diagnosis of mature T-ce neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) for thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen	
Recognizes a 67kDa transmembrane protein, which is identified as CD5 CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell are Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplast cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CD5 aberrant expression is useful in making a diagnosis of mature T-center neoplasms. Anti-CD5 detection is diagnostic in CLL/SLL within a panel of B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also useful in differentiating among mature small lymphoid cell malignancies addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) from thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.  Immunogen  Recombinant fragment (around aa450-495) of human CD5 protein (intra	bbit
CD5 antigen is found on 95% of thymocytes and 72% of peripheral bloo lymphocytes. In lymph nodes, the main reactivity is observed in T cell are Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplast cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (Comantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymphocytic lymphoma, and a subset (~10%) of diffuse large B-cell lymphocytic lymphoma, and a subset (~10%) of diffuse large B-cell lymphocytic lymphocytic leukemia/small lymphocytic lymphoma (TD5 aberrant expression is useful in making a diagnosis of mature T-center in the subset of the su	
	I eas. c B- LL/SLL), oma. f other very
	ellular
Notes  DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subside	iaries.
Product Application Details	
Applications Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffir CyTOF-ready	
Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffir CyTOF-ready	





### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NBP2-34593C**

NBP1-43319C Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 650]

NBP2-38509PEP CD5 Recombinant Protein Antigen

6507-IL-010/CF IL-4 [Unconjugated]

1636-CD-050 CD5

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34593C

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

