Product Datasheet

TfR (Transferrin R) Antibody (DF1513) [PE/Cy5.5] NBP2-34602PECY55

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34602PECY55

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34602PECY55



NBP2-34602PECY55

TfR (Transferrin R) Antibody (DF1513) [PE/Cy5.5]

TfR (Transferrin R) Antibody (DF1513) [PE/Cy5.5]	
Product Information	
0.1 ml	
Please see the vial label for concentration. If unlisted please contact technical services.	
Store at 4C in the dark. Do not freeze.	
Monoclonal	
DF1513	
0.05% Sodium Azide	
IgG1 Kappa	
PE/Cy5.5	
Protein A or G purified	
PBS	
Mouse	
7037	
TFRC	
Human	
It recognizes a ~90-95kDa protein which is identified as cell surface transferrin receptor (CD71), a disulfide-bonded homodimeric glycoprotein of 180-190kDa (Workshop IV). This monoclonal antibody is highly specific to CD71 and shows no cross-reaction with other related proteins. Ligand for transferrin receptor is the serum iron transport protein, transferrin. This receptor is broadly distributed in carcinomas, sarcomas, leukemias, and lymphomas. CD71/Transferrin receptor has been reported to be associated with cell proliferation in both normal and neoplastic tissues and useful in predicting clinical behavior or response to therapy in a number of malignancies including breast cancer.	
KG1 acute myeloid leukemia cell line	
Western Blot, Simple Western, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready	
Western Blot, Simple Western, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready	
Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any questions.	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34602PECY55

NB100-92243PEP TfR (Transferrin R) Antibody Blocking Peptide

210-TA-005 TNF-alpha [Unconjugated]

2474-TR-050 TfR (Transferrin R) [Unconjugated]

202-IL-010 IL-2 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34602PECY55

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

