Product Datasheet

Chorionic Gonadotropin beta Chain (hCG beta) Antibody (HCGb/459) [Alexa Fluor® 488] NBP2-34631AF488

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34631AF488

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34631AF488

NBP2-34631AF488

Chorionic Gonadotropin beta Chain (hCG beta) Antibody (HCGb/459) [Alexa Fluor® 488]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneHCGb/459Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1082Gene ID1082Gene SymbolCGB3Specificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but tises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chain designed as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle attimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibudy is denoted a turnors of trophoblastic origin such as choricocarionma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinoma and adenocarcinoma of the lung demonstrate antip-hCg positivity in 90% and 60% of cases respectively. 20% of ulng squamous cell carcinoma and adenocarcinoma of the lung thermostrate anti-hCg positivity in 90% and 60% of cases respectively. 20% of ulng squamous cell carcinoma and adenocarcinoma of the lung thermostrate anti-hCg positivity in 90% and 60% of cases respectively.		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneHCGb/459Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1082Gene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a dycoprotein, which is secreted in large quantiles by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunits i identical to that of thyroid stimulating hormone (LH), HCG monoclonal antibody etects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneHCGb/459Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1082HostMouseGene ID1082SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpeciesJuin on Long on the secret with the apha sub-unit, HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicie stimulating hormone (FSH), and luteinizing hormone (TSH) follicie stimulating hormone (FSH), and luteinizing hormone (TSH) follicie stimulating hormone (FSH) and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choricorcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinoma are positive. HCG expression by non-tr	Unit Size	0.1 ml
ClonalityMonoclonalCloneHCGb/459Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantifies by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is coriposed of two non-identical, non-covalently linked polypetide chains designated as the subunit is cleatical oth and thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choricarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinoma are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Concentration	
CloneHCGb/459Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1082HostMouseGene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is corposed of two non-identical, non-covalently linked polypeptide chains designated as the subuli is identical, non-covalently linked polypeptide chains design and be schoricocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic turnors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1082HostMouseGene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and Iuteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantifies by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and thurors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Clone	HCGb/459
ConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantifies by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinoma are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Isotype	IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- uit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Conjugate	Alexa Fluor 488
Product DescriptionHostMouseGene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Purity	Protein A or G purified
HostMouseGene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Buffer	50mM Sodium Borate
Gene ID1082Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Product Description	
Gene SymbolCGB3SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Host	Mouse
SpeciesHumanMarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Gene ID	1082
MarkerPregnancy & Choriocarcinoma MarkerSpecificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Gene Symbol	CGB3
Specificity/SensitivityThis monoclonal antibody reacts with a protein of 22kDa, identified as beta sub- unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.ImmunogenRecombinant full-length Chorionic Gonadotropin beta Chain (hCG beta) protein	Species	Human
 unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior. Immunogen 	Marker	Pregnancy & Choriocarcinoma Marker
		unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the subunit is identical to that of thyroid stimulating hormone (TSH) follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG monoclonal antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90% and 60% of cases respectively. 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.
	Immunogen	

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-34631AF488

IC002G	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 488]
NBP2-59545-10ug	Recombinant Human Chorionic Gonadotropin beta Chain (hCG beta) His Protein
314-BP-010	BMP-4 [Unconjugated]
DY9034-05	Chorionic Gonadotropin beta Chain (hCG beta) [Biotin]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34631AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

