Product Datasheet

Actin (Muscle Specific) Antibody (HHF35) - Azide and BSA Free NBP2-34634-0.1mg

Unit Size: 0.1 mg

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34634

Updated 4/13/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34634



NBP2-34634-0.1mg

A ST. (MA A ST. O. S. M. ST. A L. (HILFOS). A ST. A ST	
Actin (Muscle Specific) Antibody (HHF35) - Azide and BSA Free	
Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	HHF35
Preservative	No Preservative
Isotype	IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS
Product Description	
Description	1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-34230). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	58
Gene Symbol	ACTA1
Species	Human, Mouse, Rat, Canine, Chicken, Feline, Rabbit
Marker	Muscle Cell Marker
Specificity/Sensitivity	This antibody recognizes actin of skeletal, cardiac, and smooth muscle cells. It is not reactive with other mesenchymal cells except for myoepithelium. Actin can be resolved on the basis of its isoelectric points into three distinctive components: alpha, beta, and gamma in order of increasing isoelectric point. Anti-muscle specific actin recognizes alpha and gamma isotypes of all muscle groups. Non-muscle cells such as vascular endothelial cells and connective tissues are non-reactive. Also, neoplastic cells of non-muscle-derived tissue such as carcinomas, melanomas, and lymphomas are negative. It stains tumors of smooth muscle (leiomyomas and leiomyosarcomas) as well as skeletal muscle (rhabdomyomas and rhabdomyosarcomas).
Immunogen	SDS extract of human myocardium.
Product Application Details	
Applications	Western Blot, Simple Western, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, CyTOF-ready
Recommended Dilutions	Western Blot 0.5-1.0ug/ml, Simple Western, Flow Cytometry 0.5-1ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 0.5-1ug/ml, Immunohistochemistry-Paraffin 0.5-1.0ug/ml, Immunohistochemistry-Frozen 0.5-1.0ug/ml, CyTOF-ready



Application Notes

Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.

Optimal dilution for a specific application should be determined. Immunohistochemistry (No special pretreatment is required for the IHC staining of formalin-fixed, paraffin-embedded tissues.).


In Simple Western only 10 - 15 ul of the recommended dilution is used per data point. Separated by Size.

Images


Western Blot: Actin (Muscle Specific) Antibody (HHF35) - Azide and BSA Free [NBP2-34634] - Western Blot Analysis of Endometrium muscle tissue lysate using pan Actin Antibody (HHF35).




Immunohistochemistry-Paraffin: Actin (Muscle Specific) Antibody (HHF35) - Azide and BSA Free [NBP2-34634] - Formalin-fixed, paraffinembedded human Rhabdomyosarcoma stained with Actin (Muscle Specific) Antibody (HHF35).




Western Blot: Actin (Muscle Specific) Antibody (HHF35) - Azide and BSA Free [NBP2-34634] - Western Blot Analysis of Skeletal muscle tissue lysate using pan Actin Antibody (HHF35).



Immunohistochemistry-Paraffin: Actin (Muscle Specific) Antibody (HHF35) - Azide and BSA Free [NBP2-34634] - Formalin-fixed, paraffinembedded human leiomyosarcoma (10X) stained with muscle specific actin MAb (HHF35)



Simple Western: Actin (Muscle Specific) Antibody (HHF35) - Azide and BSA Free [NBP2-34634] - Electropherogram image of the corresponding Simple Western lane. pan Actin antibody was used at 10 ug/ml dilution of Muscle (Sk) lysates(s) respectively.



Simple Western: Actin (Muscle Specific) Antibody (HHF35) - Azide and BSA Free [NBP2-34634] - Simple Western lane view shows a specific band for pan Actin in 0.2 mg/ml of Muscle (Sk) lysate(s). This experiment was performed under reducing conditions using the 12-230 kDa separation system.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34634-0.1mg

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

664-LI-025 LIGHT/TNFSF14 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34634

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

