Product Datasheet

CD79A Antibody (JCB117) [Alexa Fluor® 405] NBP2-34652AF405

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34652AF405

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34652AF405



NBP2-34652AF405

CD79A Antibody (JCB117) [Alexa Fluor® 405]

Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone JCB117 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 973 Gene Symbol CD79A Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulin constitute the B cell Agr receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is generally used to complexent mit-CD20 sepecially for mature B-cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Acti-CD79a is generally used to complement anti-CD20. This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic hymphomas at is anti-CD20. This antibody will stain many of the same lymphomas as anti-CD20. This antibody will stain many of the same lymphomas and end scasionally some types of endothelial cells as well.		
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone JCB117 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on stills appe are rist in a preams at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursors B cell type, in B cell lines. Set ilymphomas, and in some lymphoma/eukemia than is anti-CD20. Attr-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as evell.	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone JCB117 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins con B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/aeukemia than is anti-CD20. but also is more likely to stain B-lymphoblastic lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas as anti-CD20, but also is more likely to stain B-lymphobla	Unit Size	0.1 ml
Clonality Monoclonal Clone JCB117 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins of precursor B cell type, in B cell lines, B cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20. Anti-CD79a is ond in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is is generally used to complement anti-CD20. Anti-CD79a is ond in the same lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomafeukemia than is anti-CD20. Anti-CD79a is outherlial cells as well. Immunogen A synthetic peptide c	Concentration	
CloneJCB117Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID973Gene SymbolCD79ASpeciesHumanMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins or B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins or B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins or B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins or B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins or B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins or B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins or B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins or B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins or B cell Stage receptor. CD79a lifs appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20. Anti-CD79a also stains more cases of plasma cell myelomas as anti-CD20. Anti-CD79a lass otains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to a	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. Ut also is more likely to stain B-lymphoblastic hymphomas and cell myeloma and occasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID973Gene SymbolCD79ASpeciesHumanMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Clone	JCB117
ConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID973Gene SymbolCD79ASpeciesHumanMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex or mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex or mb-1 and B29 polypeptides and immunoglobulins on B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines, Anti-CD79a is generally uused to complement anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas acell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomaleukemia than is anti-CD20. but also is more likely to stain B-lymphoblastic lymphomaleukemia than is anti-CD20. but also is so dia stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Isotype	IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID973Gene SymbolCD79ASpeciesHumanMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines, B cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Conjugate	Alexa Fluor 405
Product Description Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Purity	Protein A or G purified
HostMouseGene ID973Gene SymbolCD79ASpeciesHumanMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Buffer	50mM Sodium Borate
Gene ID973Gene SymbolCD79ASpeciesHumanMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Product Description	
Gene SymbolCD79ASpeciesHumanMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Host	Mouse
SpeciesHumanMarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Gene ID	973
MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Gene Symbol	CD79A
Specificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. A cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	Species	Human
 CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. Immunogen 	Marker	B-Cell Marker
	Specificity/Sensitivity	CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of
	Immunogen	

www.novusbio.com



	5 1
Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-34652AF405

IC002V	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 405]
NBP2-60209-50ug	Recombinant Human CD79A His Protein
7268-CT-100	CTLA-4 [Unconjugated]
9685-CD-050	CD79A [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34652AF405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

