Product Datasheet

CD79A Antibody (JCB117) [DyLight 650] NBP2-34652C

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34652C

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34652C



NBP2-34652C

CD79A Antibody (JCB117) [DyLight 650]

Storage Storage Store at 4C in the dark. Clonality Monoclonal Clone JCB117 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and E29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre 1 cell stage, early in maturation, and persists until the plasma cell stage where if found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. B cell lymphomas after reatment with Rituximab (anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic pepticle corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immuno	CD/9A Antibody (JCB117) [D	ylight 650]
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone JCB117 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the mority of save where it found as an intracellular component. CD79a is found in the mority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in sc myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. Anti-CD79a is generally used to complement with Rituximab (anti-CD20). This antibody will stain many of the sar lymphomas and cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoptochemistry, Immu	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone JCB117 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where i found as an intracellular component. CD79 as is found in the majority of acute leukemias of precursor B cell type, in B cell lines, Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Kiurabb (anti-CD20). This intrody will stim playma for the anti-complement anti-CD20 specially for mature B-cell lymphomas after treatment with Kiurabb (anti-CD20). This intrody will stim plaympothasic lymphomase as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomase as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomase as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomase as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomase as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomase as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomase as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomase as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomase as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomase as anti-CD20, but	Unit Size	0.1 ml
Clone JCB117 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre cell stage, early in maturation, and persists until the plasma cell stage where in found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. but also is more likely to stain B-lymphomas atter treatment with Rituximato (anti-CD20). This anti-CD20 as stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunoflicorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry, Immuno	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Clone JCB117 Preservative JCM Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer SomM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specifficity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex is until the plasm cell stage where if ound as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. B cell lymphomas atter treatment with Rituximab (anti-CD20). This anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas atter treatment with Rituximab (anti-CD20). This anti-CD79a is generally used to complement anti-CD20, but also is more likely to stain B-lympholastic lymphomas as anti-CD20, but also is more likely to stain B-lymphomas atter treatment with Rituximab (anti-CD20). This anti-CD70 as so stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytoche	Storage	Store at 4C in the dark.
Preservative 1,000	Clonality	Monoclonal
IgG1 Kappa	Clone	JCB117
Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at prefectle stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is coll lymphomas, and in sc myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphomas as anti-CD20, but also is more likely to stain B-lympholastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloima and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry,	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulin constitute the B cell Ag receptor. CD79a lifts appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where i found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituxima (anti-CD20). This antibox will stain many of the sar lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunofistochemistry, Immunofistochemistry, Immunocytochemistry, Immunofistochemistry, Immunofistochemistry, Immunofistochemistry, Immunocytochemistry, Immunocytochemistry, Immunofistochemistry, Immunofistochemistry, Immunocytochemistry, Immunocytochemistry	Isotype	IgG1 Kappa
Product Description	Conjugate	DyLight 650
Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where if found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunooflotochemistry, Immunoptochemistry, Immunoptochemistry.	Purity	Protein A or G purified
This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at prefectle stage, early in maturation, and persists until the plasma cell stage where in found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in somyelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry.	Buffer	50mM Sodium Borate
volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Mouse Gene ID 973 Gene Symbol CD79A Species Human Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry.	Product Description	
Gene Symbol CD79A Species Human B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at prefectled to cell stage, early in maturation, and persists until the plasma cell stage where if ound as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sare lymphoma/leukemia than is anti-CD20. But also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry.	Description	This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Species	Host	Mouse
Specificity/Sensitivity B-Cell Marker	Gene ID	973
B-Cell Marker	Gene Symbol	CD79A
A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where i found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well lmmunogen Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemist	Species	Human
CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre It cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well munogen Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistr	Marker	B-Cell Marker
human CD79a protein. Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry,	Specificity/Sensitivity	CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry, Immunocy	Immunogen	
Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry, Immunocytochem	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry,	Product Application Details	
	Applications	
	Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34652C

NBP1-43319C Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 650]

NBP2-60209-50ug Recombinant Human CD79A His Protein

7268-CT-100 CTLA-4 [Unconjugated] 9685-CD-050 CD79A [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34652C

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

