# **Product Datasheet**

# CD79A Antibody (JCB117 + HM47/A9) [Janelia Fluor® 549] NBP2-34653JF549

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34653JF549

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34653JF549



# NBP2-34653JF549

CD79A Antibody (JCB117 + HM47/A9) [Janelia Fluor® 549]

Storage Store at 4C in the dark.  Clonality Monoclonal  Clone JCB117 + HIM47/A9  Preservative 0.05% Sodium Azide  Isotype IgG1 Kappa/IgG1 Kappa  Conjugate Janelia Fluor 549  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines, B cell lymphomas, after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphoma/leukemia than is anti-CD200 especially for sains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Unit Size  Concentration  Please see the vial label for concentration. If unlisted please contact technical services.  Storage  Store at 4C in the dark.  Clonality  Monoclonal  Clone  JCB117 + HM47/A9  Preservative  0.05% Sodium Azide  Isotype  IgG1 Kappa/IgG1 Kappa  Conjugate  Janelia Fluor 549  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  973  Gene Symbol  CD79A  Species  Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Concentration  Please see the vial label for concentration. If unlisted please contact technical services.  Storage  Store at 4C in the dark.  Clonality  Monoclonal  Clone  JCB117 + HM47/A9  Preservative  0.05% Sodium Azide  Isotype  IgG1 Kappa/IgG1 Kappa  Conjugate  Janelia Fluor 549  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  973  Gene Symbol  CD79A  Species  Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker  B Cell Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Concentration Please see the vial label for concentration. If unlisted please contact technical services.  Storage Store at 4C in the dark.  Clonality Monoclonal Clone JCB117 + HM47/A9  Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG1 Kappa Conjugate Janelia Fluor 549  Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Mouse Gene ID 973 Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Storage Store at 4C in the dark.  Clonality Monoclonal  Clone JCB117 + HM47/A9  Preservative 0.05% Sodium Azide  Isotype IgG1 Kappa/IgG1 Kappa  Conjugate Janelia Fluor 549  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphoma/leukemia than is anti-CD200 but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD200. This antibody will stain many of the sam lymphoma/leukemia than is anti-CD200. This antibody sains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Storage Store at 4C in the dark.  Clonality Monoclonal  Clone JCB117 + HM47/A9  Preservative 0.05% Sodium Azide  Isotype IgG1 Kappa/IgG1 Kappa  Conjugate Janelia Fluor 549  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity Anonocome and B29 polypeptides and B29 polypeptides and
Clone  Clone  JCB117 + HM47/A9  Preservative  0.05% Sodium Azide  Isotype  IgG1 Kappa/IgG1 Kappa  Conjugate  Janelia Fluor 549  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  973  Gene Symbol  CD79A  Species  Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker  B Cell Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Clone JCB117 + HM47/A9  Preservative 0.05% Sodium Azide  Isotype IgG1 Kappa/IgG1 Kappa  Conjugate Janelia Fluor 549  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Clone  JCB117 + HM47/A9  Preservative  0.05% Sodium Azide  Isotype  IgG1 Kappa/IgG1 Kappa  Conjugate  Janelia Fluor 549  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  973  Gene Symbol  CD79A  Species  Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker  B Cell Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where if found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Clone  JCB117 + HM47/A9  Preservative  0.05% Sodium Azide  Isotype  IgG1 Kappa/IgG1 Kappa  Conjugate  Janelia Fluor 549  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  973  Gene Symbol  CD79A  Species  Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker  B Cell Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Preservative  Isotype  IgG1 Kappa/IgG1 Kappa Conjugate  Janelia Fluor 549  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse Gene ID  973 Gene Symbol  CD79A  Species  Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker  B Cell Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Preservative 0.05% Sodium Azide  Isotype IgG1 Kappa/IgG1 Kappa  Conjugate Janelia Fluor 549  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
IgG1 Kappa/IgG1 Kappa	Isotype IgG1 Kappa/IgG1 Kappa  Conjugate Janelia Fluor 549  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Conjugate  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  973  Gene Symbol  CD79A  Species  Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker  B Cell Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Conjugate  Purity Protein A or G purified  Buffer  50mM Sodium Borate  Product Description Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Buffer   50mM Sodium Borate	Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Product Description  Host  Mouse  Gene ID  973  Gene Symbol  Species  Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker  B Cell Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Product Description  Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Host Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Host Mouse  Gene ID 973  Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphoma'leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Gene Symbol CD79A  Species Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker B Cell Marker  Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Gene Symbol  CD79A  Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker  B Cell Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Gene Symbol  Species  Human, Mouse, Rat, Porcine, Bovine, Monkey  Marker  B Cell Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Species	Species  Human, Mouse, Rat, Porcine, Bovine, Monkey  B Cell Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Marker  Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Marker  B Cell Marker  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	Specificity/Sensitivity  A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre E cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.	CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and
Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in sor myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic
human CD79A protein (JCB117 and HM47/A9). (Uniprot: P11912)	Immunogen  A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein (ICB117 and HM47/A9) (Uniprot: P11912)
Notes Sold under license from the Howard Hughes Medical Institute, Janelia Researd Campus.	numum obrox protein (obb117 and 111014777.to). (omprot. 1 11312)
	Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research
Product Application Details	Notes Sold under license from the Howard Hughes Medical Institute, Janelia Researd Campus.
Product Application Details  Applications  Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready	Notes  Sold under license from the Howard Hughes Medical Institute, Janelia Researd Campus.  Product Application Details Applications  Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,
Applications  Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready	Notes  Sold under license from the Howard Hughes Medical Institute, Janelia Researd Campus.  Product Application Details  Applications  Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready  Recommended Dilutions  Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/
mmunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of	leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in sor myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sam lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of
· ·	ridinal Obton placin (Obt 17 and 1104-1770). (Oniplac: 1 11012)
· ·	
· ·	
· ·	
Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research	human CD79A protein (ICR117 and HM47/A9) (Uniprot: P11912)
Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research	





### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## Products Related to NBP2-34653JF549

NBP2-60209-50ug Recombinant Human CD79A His Protein

7268-CT-100 CTLA-4 [Unconjugated] 9685-CD-050 CD79A [Unconjugated]

AF114 CD45 Antibody [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34653JF549

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

