Product Datasheet

Melan-A/MART-1 Antibody (M2-7C10 + M2-9E3) [Alexa Fluor® 405] NBP2-34680AF405

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34680AF405

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34680AF405

NBP2-34680AF405

Melan-A/MART-1 Antibody (M2-7C10 + M2-9E3) [Alexa Fluor® 405]

Product Information	
---------------------	--

Unit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneM2-7C10 + M2-9E3Preservative0.05% Sodium AzideIsotypeIgG2b Lambda/IgG2b KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description2315Gene 1D2315Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma ansociated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanom	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneM2-7C10 + M2-9E3Preservative0.05% Sodium AzideIsotypeIgG2b Lambda/IgG2b KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description2315HostMouseGene ID2315Gene SymbolHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated by autologous cytotoxic nuologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosmes and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length MMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Unit Size	0.1 ml
ClonalityMonoclonalCloneM2-7C10 + M2-9E3Preservative0.05% Sodium AzideIsotypeIgG2b Lambda/IgG2b KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description4000HostMouseGene ID2315Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognized a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This monoclonal antibory labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Concentration	•
CloneM2-7C10 + M2-9E3Preservative0.05% Sodium AzideIsotypeIgG2b Lambda/IgG2b KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID2315Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognized a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified marker antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens ecognized by autologous cytotoxic T lymphocytes and endoplasmic reticulum. This monochand antibody labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelia, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG2b Lambda/IgG2b KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2315Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocyte. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocyte. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocyte. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocyte. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocyte. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocyte. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocyte. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Clonality	Monoclonal
IsotypeIgG2b Lambda/IgG2b KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2315Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityMelanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1, and GAGE-1, Ubucellular fractionation shows that MART-1 is resent in melanosomes and endoplasmic reticulum. This monoclonal antibody labels melanomas and other tumors showing mel	Clone	M2-7C10 + M2-9E3
ConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2315Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T ells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This monoclonal antibody labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat Reactivity Notes Use in Mouse reported in scientific literature (PMID:33857287) Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This monoclonal antibody labels melanomas and other tumors showing melancoytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Isotype	IgG2b Lambda/IgG2b Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2315Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Conjugate	Alexa Fluor 405
Product DescriptionHostMouseGene ID2315Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Purity	Protein A or G purified
HostMouseGene ID2315Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosames and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Buffer	50mM Sodium Borate
Gene ID2315Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This monoclonal antibody labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Product Description	
Gene SymbolMLANASpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This monoclonal antibody labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Host	Mouse
SpeciesHuman, Mouse, RatReactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This monoclonal antibody labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Gene ID	2315
Reactivity NotesUse in Mouse reported in scientific literature (PMID:33857287)MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Gene Symbol	MLANA
MarkerMelanoma MarkerSpecificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This monoclonal antibody labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Species	Human, Mouse, Rat
Specificity/SensitivityThis antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This monoclonal antibody labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.ImmunogenRecombinant full-length hMelan-A/MART-1 protein (M2-7C10; M2-9E3) (Uniprot:	Reactivity Notes	Use in Mouse reported in scientific literature (PMID:33857287)
 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This monoclonal antibody labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen 	Marker	Melanoma Marker
\bullet	Specificity/Sensitivity	(Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This monoclonal antibody labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.
	Immunogen	



Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Western Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready
Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Flow (Intracellular), CyTOF-ready





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-34680AF405

NBP2-23223	Recombinant Human Melan-A/MART-1 His Protein
285-IF-100	IFN-gamma [Unconjugated]
NBP2-33535PEP	Melan-A/MART-1 Recombinant Protein Antigen
202-IL-010	IL-2 [Unconjugated]
202 .2 0 .0	[0

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34680AF405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

