Product Datasheet

CD63 Antibody (MX-49.129.5) [DyLight 680] NBP2-34689FR

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34689FR

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34689FR



NBP2-34689FR

CD63 Antibody (MX-49.129.5) [DyLight 680]

as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Full length CD63 of human origin (Uniprot: P08962) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	OD03 / (III)D0dy (W/X 43.123.3) [DyEight 000]		
Please see the vial label for concentration. If unlisted please contact technical services.	Product Information		
Storage Store at 4C in the dark. Clonality Monoclonal Clone MX-49,129.5 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP/35/29. The tetraspanias are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspania re integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspania re integral membrane and pranulocytes, To cell and B cells. It is located on the basophilic granule membranes on a membrane and macrophages, and is weakly expressed on granulocytes, To cell and B cells. It is located on the basophilic granule membrane and the plasma membrane of lymphocytes and granulocytes. CD63 as a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD63, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Full length CD63 of human origin (Uniprot: P08962) Product Application Details Applications	Unit Size	0.1 ml	
Clone MX-49.129.5 Preservative 0.05% Sodium Azide lsotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Full length CD63 of human origin (Uniprot: P08962) Product Application Details Applications	Concentration	·	
Clone MX-49.129.5 Preservative 0.05% Sodium Azide Isotype 1gG1 Kappa DyLight 680 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/629. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to he plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, To cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a membrane membrane of the TM4 superfamily of leukocyte glycoprotein stat includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 is a membrane of the TM4 superfamily of leukocyte glycoprotein stat includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 is a membrane of the TM4 superfamily of leukocyte glycoprotein stat includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 is a membrane of the TM4 superfamily of leukocyte glycoprotein stat includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 is a membrane of the TM4 superfamily of leukocyte glycoprotein stat includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 is a membrane of the TM4 superfamily of leukocyte glycoprotein stat includes CD9, CD37 and CD53, which contain four tran	Storage	Store at 4C in the dark.	
Preservative 1,05% Sodium Azide	Clonality	Monoclonal	
IgG1 Kappa	Clone	MX-49.129.5	
Conjugate DyLight 680 Purity Protein A purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Preservative	0.05% Sodium Azide	
Purity Protein A purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes. CD63 is a membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes. To cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Full length CD63 of human origin (Uniprot: P08962) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Isotype	IgG1 Kappa	
Buffer 50mM Sodium Borate	Conjugate	DyLight 680	
Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Full length CD63 of human origin (Uniprot: P08962) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Purity	Protein A purified	
This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Full length CD63 of human origin (Uniprot: P08962) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Buffer	50mM Sodium Borate	
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Full length CD63 of human origin (Uniprot: P08962) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Product Description		
Gene Symbol CD63 Species Human, Mouse Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Full length CD63 of human origin (Uniprot: P08962) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Description	volume of this product. The volume will be greater than or equal to the unit size	
Species	Host	Mouse	
Species	Gene ID	967	
MarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.ImmunogenFull length CD63 of human origin (Uniprot: P08962)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsApplicationsApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Gene Symbol	CD63	
This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Full length CD63 of human origin (Uniprot: P08962) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Species	Human, Mouse	
as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Full length CD63 of human origin (Uniprot: P08962) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Marker	Late Endosomes Marker	
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Specificity/Sensitivity	The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during	
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Immunogen	Full length CD63 of human origin (Uniprot: P08962)	
Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	
	Product Application Details		
,, , . , . , ,	Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready	



Recommended Dilutions

Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722

Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34689FR

NBP1-43319FR-0.5ml Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 680]

H00000967-G01-2ug Recombinant Human CD63 Protein

203-IL-010 IL-3 [Unconjugated] 5048-CD-050 CD63 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34689FR

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

