Product Datasheet

CD45RB Antibody (PD7/26) [Alexa Fluor® 350] NBP2-34708AF350

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34708AF350

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34708AF350



NBP2-34708AF350

CD45RB Antibody (PD7/26) [Alexa Fluor® 350]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClonePD7/26Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBufferSomM Sodium BorateProduct Description5788Gene ID5788Gene SymbolPTPRCSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Mutiple isoforms of CD45R are distributed for CD45RA (exon 4), CD45RR (exon son 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 Spliced out), The variation in these isoforms is localized to the extracelluar domain of CD45R, while the initracelluar domain is conserved. CD45RB (exon 4), CD45RR (exon son 5), CD45RC (exon 5) and CD45RD (exons 4 to 5 Spliced out), The variation in these isoforms is localized to the extracelluar domain of CD45R, while the initracelluar domain is conserved. CD45RB (exon 4), CD45RR (exon son 5), CD45RC (exon 6) and CD45RD (exon 4), CD45RR (exon son 5), CD45RC (exon 6) and CD45RD (exon 4), CD45RR (exon son 5), CD45RC (exon 6) and CD45RD (exon 4), CD45RR (exon son 5), CD45RC (exon 6) and CD45RD (exon 4), CD45RR (exon son 5), CD45RC (exon 6) and CD45RD (exon 4), CD45RR (exon son 5), CD45RC (exon 6) and CD45RD (exon 4), CD45RR (exon son 5), CD45RC (exon 6) and CD45RD (exon 4) to CD45RA (exon 4), CD45RR (exon son 6) in the extracelluar domain is conserved. CD45RB is expressed on mature B- tymphocytes and the major	, -		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClonePD7/26Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description5788Gene ID5788Gene SymbolPTPRCSpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of moncclonal antibodies specific for CD45R, while the initracellular domain is conserved. CD45R Bi expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Product Information		
services.StorageStore at 4C in the dark.ClonalityMonoclonalClonePD7/26Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1000000000000000000000000000000000000	Unit Size	0.1 ml	
ClonalityMonoclonalClonalityMonoclonalClonePD7/26Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanSpeciesCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RA (exon 5), CD45RB (exon 5), CD45RC (exon 6) and CD45RA (exon 4) CD45RB (exon 4), CD45RB (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RA (exon 4) for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RA (exon 4) for CD45RB (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RA (exon 4) for CD45RB (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RA (exon 4) for CD45RB (exon 5), CD45RC (exon 6) and CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Concentration		
ClonePD7/26Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RD (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Storage	Store at 4C in the dark.	
Preservative0.05% Sodium AzidePreservativelgG1 KappaIsotypelgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID5788Gene SymbolPTPRCSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific or CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Clonality	Monoclonal	
IsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Clone	PD7/26	
ConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Preservative	0.05% Sodium Azide	
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Isotype	IgG1 Kappa	
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B- lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Conjugate	Alexa Fluor 350	
Product DescriptionHostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B- lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Purity	Protein A or G purified	
HostMouseGene ID5788Gene SymbolPTPRCSpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B- lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Buffer	50mM Sodium Borate	
Gene ID5788Gene SymbolPTPRCSpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B- lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Product Description		
Gene SymbolPTPRCSpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B- lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Host	Mouse	
SpeciesHumanSpecificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B- lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Gene ID	5788	
Specificity/SensitivityCD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B- lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Gene Symbol	PTPRC	
transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B- lymphocytes and the majority of lymphomas and leukemias of B-cell origin.	Species	Human	
Immunogen Human peripheral blood lymphocytes maintained in T cell growth factor	Specificity/Sensitivity	transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-	
	Immunogen	Human peripheral blood lymphocytes maintained in T cell growth factor	



Notes	

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details

	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready
	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Images

CD45RB Antibody (PD7/26) [Alexa Fluor® 350] [NBP2-34708AF350] - Vial of Alexa Fluor 350 conjugated antibody. Alexa Fluor 350 is optimally excited at 346 nm by the UV laser (350 or 355 nm) and has an emission maximum of 442 nm.

www.novusbio.com

Alexa Fluor [®] 350				
Alexa Fluor ^e 5 ⁶⁰ EXCITATION MAX (nm) EMISSION MAX (nm)				
Alleza Filuor ^e 3 ⁵⁰ UV (350) 450/45 EXCITATION MAX (nm) EMISSION MAX (nm)		Alexa Fluc	or® 350	
EXCITATION MAX (mm) EMISSION MAX (nm)		LASER (nm)	FILTER]
	Alexa Fluor® 350	UV (350)	450/45	
346 442		EXCITATION MAX (nm)	EMISSION MAX (nm)	1
		346	442	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-34708AF350

6507-IL-010/CF	IL-4 [Unconjugated]
M6000B-1	IL-6 [HRP]
10477-CD-050	CD45RB [Unconjugated]
210-TA-005	TNF-alpha [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34708AF350

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

