Product Datasheet

CD34 Antibody (SPM123) [DyLight 488] NBP2-34733G

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34733G

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34733G



NBP2-34733G

CD34 Antibody (SPM123) [DyLight 488]

Product Information	, , , , , ,	•
Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark.	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM123 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes on monoclonal antibodies to CD34 are classified into three main categories, class I, and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopanic. CD34 expression is a hallmark for identifying pluripotent hemain categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopanic. CD34 expression is a hallmark for identifying pluripotent hemain categories, class I, class II san class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopanic. CD34 expression is a hallmark for identifying pluripotent hemain categories class II, it is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopanic. CD34 expressed by septifier antiae cD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft fisset utmorited progenitors differentiae. CD34 is a marker of choice for st	Unit Size	0.1 ml
Clone SPM123 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differential sensitivity to degradation by specific enzymes committed progenitors differentiales c.D34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the no	Concentration	•
Clone SPM123 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class II, alt is a class III antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiale. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express than molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express the molecule more than the non-proli	Storage	Store at 4C in the dark.
Preservative	Clonality	Monoclonal
IgG1 Kappa	Clone	SPM123
Conjugate DyLight 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into there main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapin. CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentials. CD34 is a marker of choice for staining plasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molec	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enymose, spitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitivity to glycoprotease and cymopapain. CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non	Isotype	IgG1 Kappa
Buffer 50mM Sodium Borate	Conjugate	DyLight 488
Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 947 Gene Symbol Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Purity	Protein A or G purified
This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endoth	Buffer	50mM Sodium Borate
wolume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta	Product Description	
Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Description	volume of this product. The volume will be greater than or equal to the unit size
CD34	Host	Mouse
Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain. CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta	Gene ID	947
(Negative), Rat (Negative) Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Gene Symbol	CD34
Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Species	
This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Reactivity Notes	Does not react with Rat, Bovine or Canine.
glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Marker	Hematopoietic Stem Cell & Endothelial Marker
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Specificity/Sensitivity	glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Immunogen	Detergent solubilized vesicular suspension prepared from human term placenta
		DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

Product Application Details

Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready **Applications**



Recommended Dilutions

Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34733G

NBP1-43319G-0.5ml Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 488]

NBP2-53044-50ug Recombinant Human CD34 His Protein

210-TA-005 TNF-alpha [Unconjugated]

9655-CD-050 CD34 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34733G

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

