Product Datasheet

CD34 Antibody (SPM123) [DyLight 350] NBP2-34733UV

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34733UV

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34733UV



NBP2-34733UV

CD34 Antibody (SPM123) [DyLight 350]

Product Information	, , , , , , , , , , , , , , , ,	
Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark.	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM123 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, eithopes of monoclonal antibodies to CD34 are classified into three main categories, class I, and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapina. (CD34 expression is a hallmark for identifying pluripotent hemain categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapina. (CD34 expression is a hallmark for identifying pluripotent hemain categories, class II, a list a class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapina. (CD34 expression is a hallmark for identifying pluripotent hemain categories, class II, a list a class II and the main categories, class II, a list a class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapina. (CD34 expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for stainin	Unit Size	0.1 ml
Clone SPM123 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class II, alt is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiales. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating	Concentration	·
Clone SPM123 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34, On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class II, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. It is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelian. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-prolifer	Storage	Store at 4C in the dark.
Preservative D.0.5% Sodium Azide	Clonality	Monoclonal
IgG1 Kappa	Clone	SPM123
Conjugate DyLight 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapian. CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentials. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitivity to glycoprotease and cympopapain. CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by schorable and sacute myeloid leukemia. In addition, CD34 is expressed in its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cel	Isotype	IgG1 Kappa
Product Description	Conjugate	DyLight 350
Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 947 Gene Symbol Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proli	Purity	Protein A or G purified
This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express this molecule more than the non-profilerating endothelial cells express than the non-profilerating endothelial cells express	Buffer	50mM Sodium Borate
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 947 Gene Symbol CD34 Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelian cells express that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells express this molecule	Product Description	
Gene Symbol Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is expressed by soft itssue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene Symbol Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative) Reactivity Notes Does not react with Rat, Bovine or Canine. Marker Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain. CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Host	Mouse
Species Human, Cynomolgus Monkey, Rhesus Macaque, Bovine (Negative), Canine (Negative), Rat (Negative)	Gene ID	947
(Negative), Rat (Negative) Does not react with Rat, Bovine or Canine. Hematopoietic Stem Cell & Endothelial Marker Specificity/Sensitivity This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Gene Symbol	CD34
Marker Hematopoietic Stem Cell & Endothelial Marker This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Species	
This monoclonal antibody recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Reactivity Notes	Does not react with Rat, Bovine or Canine.
glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower specificity. Immunogen Detergent solubilized vesicular suspension prepared from human term placenta Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Marker	Hematopoietic Stem Cell & Endothelial Marker
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Specificity/Sensitivity	glycosylated protein of 90-120kDa, which is identified as CD34. On the basis of differential sensitivity to degradation by specific enzymes, epitopes of monoclonal antibodies to CD34 are classified into three main categories, class I, class II and class III. It is a class II antibody whose epitope is resistant to neuraminidase but sensitive to glycoprotease and chymopapain.CD34 expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposis sarcoma, but with a lower
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Immunogen	Detergent solubilized vesicular suspension prepared from human term placenta
	Notes Product Application Details	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

Product Application Details

Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready **Applications**



Recommended Dilutions

Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34733UV

NBP1-43319UV-0.5ml Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 350]

NBP2-53044-50ug Recombinant Human CD34 His Protein

210-TA-005 TNF-alpha [Unconjugated]

9655-CD-050 CD34 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34733UV

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

