# **Product Datasheet**

# MUC1 Antibody (SPM132) [Alexa Fluor® 405] NBP2-34737AF405

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34737AF405

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34737AF405



# NBP2-34737AF405

MUC1 Antibody (SPM132) [Alexa Fluor® 405]

Product Information Unit Size  0.1 ml  Concentration  Please see the vial label for concentration. If unlisted please contact te services.  Storage  Store at 4C in the dark.  Clonality  Monoclonal  Clone  SPM132  Preservative  0.05% Sodium Azide  Isotype  IgG1 Kappa  Conjugate  Alexa Fluor 405  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  4582  Gene Symbol  MUC1  Species  Human  Marker  Specificity/Sensitivity  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with tepitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack, subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Concentration  Please see the vial label for concentration. If unlisted please contact te services.  Storage  Store at 4C in the dark.  Clonality  Monoclonal  Clone  SPM132  Preservative  0.05% Sodium Azide  Isotype  IgG1 Kappa  Conjugate  Alexa Fluor 405  Purity  Protein A or G purified  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  4582  Gene Symbol  MUC1  Species  Human  Marker  Epithelial Marker  Specificity/Sensitivity  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with te epitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack, subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Storage Store at 4C in the dark.  Clonality Monoclonal  Clone SPM132  Preservative 0.05% Sodium Azide  Isotype IgG1 Kappa  Conjugate Alexa Fluor 405  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 4582  Gene Symbol MUC1  Species Human  Marker Epithelial Marker  Specificity/Sensitivity  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with tepitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack. subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Clonality Monoclonal Clone SPM132 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Mouse Gene ID 4582 Gene Symbol MUC1 Species Human Marker Epithelial Marker Specificity/Sensitivity In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with tepitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack. Subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Clone  SPM132  Preservative  0.05% Sodium Azide  Isotype  IgG1 Kappa  Conjugate  Alexa Fluor 405  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  4582  Gene Symbol  MUC1  Species  Human  Marker  Epithelial Marker  Specificity/Sensitivity  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with tepitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may p protective layer on epithelial cells against bacterial and enzyme attack subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Preservative  Isotype  IgG1 Kappa  Conjugate  Alexa Fluor 405  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Mouse  Gene ID  4582  Gene Symbol  MUC1  Species  Human  Marker  Epithelial Marker  Specificity/Sensitivity  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with the epitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack. Subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Isotype Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Mouse Gene ID 4582 Gene Symbol MUC1 Species Human Marker Epithelial Marker Specificity/Sensitivity In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with tepitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack, subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Conjugate Alexa Fluor 405  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 4582  Gene Symbol MUC1  Species Human  Marker Epithelial Marker  Specificity/Sensitivity In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with the epitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack. Subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description Host Mouse Gene ID 4582 Gene Symbol MUC1 Species Human Marker Epithelial Marker  Epithelial Marker  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with the epitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack, subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Buffer 50mM Sodium Borate  Product Description  Host Mouse  Gene ID 4582  Gene Symbol MUC1  Species Human  Marker Epithelial Marker  Specificity/Sensitivity In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with the epitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may pure protective layer on epithelial cells against bacterial and enzyme attack, subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Product Description  Host Mouse  Gene ID 4582  Gene Symbol MUC1  Species Human  Marker Epithelial Marker  Specificity/Sensitivity In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with the epitope in the tandem repeats. The alpha subunit has cell adhesive procant act both as an adhesion and an anti-adhesion protein. EMA may pure protective layer on epithelial cells against bacterial and enzyme attacks subunit contains a C-terminal domain, which is involved in cell signaling phosphorylations and protein-protein interactions. In immunohistochem
Host Mouse  Gene ID 4582  Gene Symbol MUC1  Species Human  Marker Epithelial Marker  Specificity/Sensitivity In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with the epitope in the tandem repeats. The alpha subunit has cell adhesive procant act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack, subunit contains a C-terminal domain, which is involved in cell signaling phosphorylations and protein-protein interactions. In immunohistochem
Gene Symbol  MUC1  Species  Human  Epithelial Marker  Specificity/Sensitivity  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with tepitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack. subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
Gene Symbol  MUC1  Species  Human  Epithelial Marker  Specificity/Sensitivity  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with the epitope in the tandem repeats. The alpha subunit has cell adhesive procaute act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack. Subunit contains a C-terminal domain, which is involved in cell signaling phosphorylations and protein-protein interactions. In immunohistochem
Species  Human  Epithelial Marker  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with the epitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attacks subunit contains a C-terminal domain, which is involved in cell signaling phosphorylations and protein-protein interactions. In immunohistochem
Marker  Specificity/Sensitivity  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with the epitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attacks subunit contains a C-terminal domain, which is involved in cell signaling phosphorylations and protein-protein interactions. In immunohistochem
Specificity/Sensitivity  In Western blotting, it recognizes proteins in MW range of 265-400kDa as different glycoforms of EMA. This monoclonal antibody reacts with tepitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack. subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
as different glycoforms of EMA. This monoclonal antibody reacts with tepitope in the tandem repeats. The alpha subunit has cell adhesive procan act both as an adhesion and an anti-adhesion protein. EMA may protective layer on epithelial cells against bacterial and enzyme attack. subunit contains a C-terminal domain, which is involved in cell signalin phosphorylations and protein-protein interactions. In immunohistochem
assays, it superbly stains routine formalin/paraffin carcinoma tissues. A EMA is useful as a pan-epithelial marker for detecting early metastatic carcinoma in bone marrow or liver.
Immunogen Human milk fat globule membranes (Uniprot: P15941)
Alexa Fluor (R) products are provided under an intellectual property lic Life Technologies Corporation. The purchase of this product conveys to buyer the non-transferable right to use the purchased product and composed of the product only in research conducted by the buyer (whether the bure academic or for-profit entity). The sale of this product is expressly concurred the buyer not using the product or its components, or any materials may the product or its components, in any activity to generate revenue, which include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (in resale, regardless of whether they are resold for use in research. For in on purchasing a license to this product for purposes other than as descrepted above, contact Life Technologies Corporation, 5791 Van Allen Way, Can 92008 USA or outlicensing@lifetech.com. This conjugate is made of demand. Actual recovery may vary from the stated volume of this product volume will be greater than or equal to the unit size stated on the dataset.



<b>Product Application Details</b>	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Products Related to NBP2-34737AF405

IC002V Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 405]

H00004582-Q01-10ug Recombinant Human MUC1 GST (N-Term) Protein

1129-ER-050 ErbB2/Her2 [Unconjugated]

10332-MU-050 MUC1 [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34737AF405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

