Product Datasheet

CD63 Antibody (SPM524) [DyLight 405] NBP2-34779V

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34779V

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34779V



NBP2-34779V

CD63 Antibody (SPM524) [DyLight 405]

Unit Size 0.1 mi Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM524 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Bymbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on elluriface and granular membrane of hematopoletic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane of hematopoletic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane of lymphocytes and granulacytes. CD63 is a membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and membranes of lymphocytes and granulocytes. CD63 is a membrane of in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hemansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Product Application Details Product Application Details	02007 minocay (01 mio2 1) [2	y=.g 100]
Please see the vial label for concentration. If unlisted please contact technical services.	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM524 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of Imphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoprotein that iranslocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of hematopomen progression. Is not transmembrane egions. CD63 may play one lein phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.	Unit Size	0.1 ml
Clone SPM524 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells Notes	Concentration	·
Clone SPM524 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoletic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on granulozies, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes and macrophages, and is weakly expressed on granulozies. T cell and B cells. It is located on the basophilic granule membranes and on the plasma membrane and macrophages, and is weakly expressed on granulozies. T cell and B cells. It is located on the basophilic granule membranes and on the plasma membrane associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Storage	Store at 4C in the dark.
Preservative IgG1 Kappa IgG1 Kappa IgG1 Kappa DyLight 405 Purity	Clonality	Monoclonal
IgG1 Kappa	Clone	SPM524
Conjugate DyLight 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glocoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membrane and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that include CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and macrophages, and is weakly expressed on activated platelets, monocytes and	Isotype	IgG1 Kappa
Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes. T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Conjugate	DyLight 405
Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 me play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Purity	Protein A or G purified
This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Buffer	50mM Sodium Borate
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 967 Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Product Description	
Gene Symbol Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene Symbol CD63 Species Human, Mouse Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Host	Mouse
Reactivity Notes Mouse reactivity reported in scientific literature (PMID: 31207235). Marker Late Endosomes Marker Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Gene ID	967
Reactivity NotesMouse reactivity reported in scientific literature (PMID: 31207235).MarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.ImmunogenSmooth plasma membrane fraction of MeWo cellsNotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Gene Symbol	CD63
MarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.ImmunogenSmooth plasma membrane fraction of MeWo cellsNotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Species	Human, Mouse
This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 31207235).
as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. Immunogen Smooth plasma membrane fraction of MeWo cells Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Marker	Late Endosomes Marker
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Specificity/Sensitivity	The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multimolecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Immunogen	Smooth plasma membrane fraction of MeWo cells

Product Application Details

Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin **Applications**



Recommended Dilutions

Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34779V

NBP1-43319V-0.5ml Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 405]

H00000967-G01-2ug Recombinant Human CD63 Protein

203-IL-010 IL-3 [Unconjugated] 5048-CD-050 CD63 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34779V

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

