Product Datasheet

CD79A Antibody (SPM550) [DyLight 680] NBP2-34792FR

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34792FR

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34792FR



NBP2-34792FR

CD79A Antibody (SPM550) [DyLight 680]

District Concentration	
Please see the vial label for concentration. If unlisted please contact ted services.	
Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM550 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B2: CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulins on B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines, Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas are miteratment with Rituximab (anti-CD20). This antibody will stain many of thymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more oal plasma cell myeloma and occasionally some types of endothelial cells a limmunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADV/human CD79A protein. (Uniprot: P11912)	
Clone SPM550 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B2t CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage w found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of tlymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more can plasma cell myeloma and occasionally some types of endothelial cells a limmunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADV/human CD79A protein. (Uniprot: P11912)	hnical
Clone SPM550 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B2t CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and E29 polypeptide immunoglobulins constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage we found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of tlymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more caplasma cell myeloma and occasionally some types of endothelial cells a limmunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVendeman CD79A protein. (Uniprot: P11912)	
Preservative Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the ustated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B2's CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage we found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines, Anti-CD79a is gen used to complement anti-CD20. This antibody will stain many of the lymphomas as anti-CD20, but also is more likely to stain B-lymphoblast lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more capiasma cell myeloma and occasionally some types of endothelial cells as Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVenduman CD79A protein. (Uniprot: P11912)	
Isotype	
Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the ustated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B22 CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persist he plasma cell stage we found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of the lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more caplasma cell myeloma and occasionally some types of endothelial cells a limmunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADV/human CD79A protein. (Uniprot: P11912)	
Purity Buffer Product Description This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the ustated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B2: CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20. This antibody will stain many of the lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more caplasma cell myeloma and occasionally some types of endothelial cells a limmunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADV/human CD79A protein. (Uniprot: P11912)	
Product Description This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the distance on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B2th CD79b) polypeptides, is non-covalently associated with membrane-bout immunoglobulin on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage we found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of the lymphomas as anti-CD20, but also is more likely to stain B-lymphoblast lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more caplasma cell myeloma and occasionally some types of endothelial cells at Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADV/human CD79A protein. (Uniprot: P11912)	
Product Description This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the distance on the datasheet. Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B2th CD79b) polypeptides, is non-covalently associated with membrane-bour immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage we found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines, Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of the lymphomas as anti-CD20. Anti-CD79a also stains more can plasma cell myeloma and occasionally some types of endothelial cells at Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADV/human CD79A protein. (Uniprot: P11912)	
This conjugate is made on demand. Actual recovery may vary from the volume of this product. The volume will be greater than or equal to the ustated on the datasheet. Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B2t CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage we found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of t lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more caplasma cell myeloma and occasionally some types of endothelial cells a lammunogen Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVI) human CD79A protein. (Uniprot: P11912)	
volume of this product. The volume will be greater than or equal to the ustated on the datasheet. Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B2t CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulins constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage we found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of to lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more caplasma cell myeloma and occasionally some types of endothelial cells a limmunogen Immunogen Notes	
Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B2t CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage we found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of the lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more caplasma cell myeloma and occasionally some types of endothelial cells at Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADV/human CD79A protein. (Uniprot: P11912)	
Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of the lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more caplasma cell myeloma and occasionally some types of endothelial cells a lammunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADV/human CD79A protein. (Uniprot: P11912)	
Species	
Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B25 CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears cell stage, early in maturation, and persists until the plasma cell stage of found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of the lymphomas as anti-CD20, but also is more likely to stain B-lymphoblast lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more can plasma cell myeloma and occasionally some types of endothelial cells at the limit of the	
Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage w found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of tlymphomas as anti-CD20, but also is more likely to stain B-lymphoblast lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more caplasma cell myeloma and occasionally some types of endothelial cells a lmmunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADV6 human CD79A protein. (Uniprot: P11912)	
CD79b) polypeptides, is non-covalently associated with membrane-bou immunoglobulins on B cells. This complex of mb-1 and B29 polypeptide immunoglobulin constitute the B cell Ag receptor. CD79a first appears a cell stage, early in maturation, and persists until the plasma cell stage w found as an intracellular component. CD79a is found in the majority of a leukemias of precursor B cell type, in B cell lines, B cell lymphomas, an myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is gen used to complement anti-CD20 especially for mature B-cell lymphomas treatment with Rituximab (anti-CD20). This antibody will stain many of t lymphomas as anti-CD20, but also is more likely to stain B-lymphoblast lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more caplasma cell myeloma and occasionally some types of endothelial cells a lmmunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVenuman CD79A protein. (Uniprot: P11912)	
human CD79A protein. (Uniprot: P11912) Notes	nd s and t pre B there it is cute d in some erally after ne same c ses of
	2) of
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsi	diaries.
Product Application Details	
Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluoresce Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready	nce,
Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry-Paraffin, CyTOF-ready	nistry/





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722

Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34792FR

NBP1-43319FR-0.5ml Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 680]

NBP2-60209-50ug Recombinant Human CD79A His Protein

7268-CT-100 CTLA-4 [Unconjugated] 9685-CD-050 CD79A [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34792FR

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

