Product Datasheet

CD79A Antibody (SPM550) [DyLight 550] NBP2-34792R

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34792R

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34792R



NBP2-34792R

CD79A Antibody (SPM550) [DyLight 550]

Storage Stora 4 C in the dark. Clonality Monoclonal Clone SPM550 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 550 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and E29 polypeptides and immunoglobulina on all persists until the plasma cell stage, early in maturation, and persists until the plasma cell stage, where found as an intracellular component. CD79a is found in the majority dacute leukemias of precursor B cell type, in B cell lines. B cell lymphomas after found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a lise penerally used to complement anti-CD20 be specially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20. This antibody will stain many of the sar lymphomas as anti-CD20 be specially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptice corresponding to as 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Western Blot, Flow Cytometry, Immunocytochemistry/ Immunoflluorescence, Immunohistochemistry, Immunoflistochemistry, Immunoflistochemistry, Immunoflistochemistry, Immunoflistochemistry, Immunoflistochemistry.	CD79A Antibody (SPIVI550) [L	Dylight 550]
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM550 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 550 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Mouse Gene ID Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where i found as an intracellular component. CD79a is found in the mority of each surplement anti-CD20. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar hymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar hymphomas and cell myelome and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to a a 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry. Immunocytochemistry	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM550 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 550 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where i found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in sc myelomas. It is not present in myelolor of T cell lines, Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Kiumabo (anti-CD20). This instrubedy will stain many of the sar lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Specially for mature B-cell lymphomas are lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries in the plasma cell myeloma and occasionality some types of endothelial cells as well immunohistochemistry, Immunocytochemistry,	Unit Size	0.1 ml
Clone SPM550 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 550 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex is until the plasma cell stage where is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas after treatment with Rituximab (anti-CD20). This anti-CD20 as tain many of the sar lymphomas as anti-CD20, but also is more likely to stain B-lymphomas atter treatment with Rituximab (anti-CD20). This anti-CD20 stain amany of the sar lymphoma allest are treatment with Rituximab (anti-CD20). This anti-CD20 stain sub stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen Immunogen A synthetic peptide corresponding to an 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemis	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Clone SPM550 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Onjugate DyLight 550 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex suntil the plasm cell stage where in cell stage, early in maturation, and persists until the plasm cell stage where found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. B cell lymphomas and in myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. but also is more likely to stain B-lymphoblastic lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas and instin-CD20. Intra-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to an 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunofistochemistry, Immunofistochemistry, Immunofistochemistry, Immunofistochemistry, Immunofistochemistry, Immunocytochemistry, Immunocytoch	Storage	Store at 4C in the dark.
Preservative 1,00% Sodium Azide	Clonality	Monoclonal
IgG1 Kappa	Clone	SPM550
Conjugate DyLight 550 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is cell lymphomas, and in sc myelomas. It is not present in myeloid or T cell lines, Anti-CD79a is generally used to complement anti-CD20. but also is more likely to stain B-lympholastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immuno	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where if ound as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituxima (anti-CD20). This antibow will stain many of the sar lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry, Immunocytochem	Isotype	IgG1 Kappa
Product Description	Conjugate	DyLight 550
Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Host Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where if ound as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas, and in st myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. Distance and single	Purity	Protein A or G purified
This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at prefectled stage, early in maturation, and persists until the plasma cell stage where found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in somyelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry.	Buffer	50mM Sodium Borate
volume of this product. The volume will be greater than or equal to the unit siz stated on the datasheet. Mouse Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where if found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry	Product Description	
Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Monkey Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where i found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry.	Description	This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Species	Host	Mouse
Specificity/Sensitivity B-Cell Marker	Gene ID	973
Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at prefectled stage, early in maturation, and persists until the plasma cell stage where it found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in somyelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well munogen Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocyto	Gene Symbol	CD79A
A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well lmmunogen Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistr	Species	Human, Mouse, Rat, Porcine, Bovine, Monkey
CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre I cell stage, early in maturation, and persists until the plasma cell stage where if found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in so myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the sar lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well mmunogen Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry,	Marker	B-Cell Marker
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry, I	Specificity/Sensitivity	CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry, Immunocy	Immunogen	
Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry, Immunocytochem	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry,	Product Application Details	
	Applications	
	Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34792R

NBP1-43319R Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 550]

NBP2-60209-50ug Recombinant Human CD79A His Protein

7268-CT-100 CTLA-4 [Unconjugated] 9685-CD-050 CD79A [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34792R

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

