Product Datasheet

Cdc20 Antibody (AR12) - IHC-Prediluted NBP2-44406

Unit Size: 7 ml

Store at 4C.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-44406

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-44406



NBP2-44406

Cdc20 Antibody (AR12) - IHC-Prediluted


services.StorageStore at 4C.ClonalityMonoclonalCloneAR12Preservative0.05% Sodium AzideIsotypeIgG1 KappaPurityProtein A or G purifiedBuffer10 mM PBS with 0.05% BSATarget Molecular Weight55 kDaProduct DescriptionThe prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining.HostMouseGene ID991Gene SymbolCDC20SpeciesHumanImmunogenUrea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834)Product Application DetailsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry (Formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.		
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C. Clonality Monoclonal Clone AR12 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 55 kDa Product Description The prediluted antibody does not require any mixing, dilution, reconstitution, of titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 991 Gene Symbol CDC20 Species Human Immunogen Urea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834) Product Application Details Applications Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry (formalin-fixed): 1-2ug/ml for 30 minutes at RT. Stain of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Information	
services.StorageStore at 4C.ClonalityMonoclonalCloneAR12Preservative0.05% Sodium AzideIsotypeIgG1 KappaPurityProtein A or G purifiedBuffer10 mM PBS with 0.05% BSATarget Molecular Weight55 kDaProduct DescriptionThe prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining.HostMouseGene ID991Gene SymbolCDC20SpeciesHumanImmunogenUrea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834)Product Application DetailsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry (Formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Unit Size	7 ml
Clonality Monoclonal Clone AR12 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 55 kDa Product Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 991 Gene Symbol CDC20 Species Human Immunogen Urea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834) Product Application Details Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Stain of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 r EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Clone AR12 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 55 kDa Product Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 991 Gene Symbol CDC20 Species Human Immunogen Urea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834) Product Application Details Applications Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Stain of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Storage	Store at 4C.
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 55 kDa Product Description The prediluted antibody does not require any mixing, dilution, reconstitution, of titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 991 Gene Symbol CDC20 Species Human Immunogen Urea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834) Product Application Details Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 rEDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clonality	Monoclonal
Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 55 kDa Product Description 55 kDa Description The prediluted antibody does not require any mixing, dilution, reconstitution, of titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 991 Gene Symbol CDC20 Species Human Immunogen Urea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834) Product Application Details Applications Application Notes Immunohistochemistry, Immunohistochemistry-Paraffin Fecommended Dilutions Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clone	AR12
Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 55 kDa Product Description Description Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 991 Gene Symbol CDC20 Species Human Immunogen Urea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834) Product Application Details Applications Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Stain of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	0.05% Sodium Azide
Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 55 kDa Product Description The prediluted antibody does not require any mixing, dilution, reconstitution, of titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 991 Gene Symbol CDC20 Species Human Immunogen Urea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834) Product Application Details Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 reDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG1 Kappa
Target Molecular Weight55 kDaProduct DescriptionThe prediluted antibody does not require any mixing, dilution, reconstitution, of titration; the antibody is ready-to-use and optimized for staining.HostMouseGene ID991Gene SymbolCDC20SpeciesHumanImmunogenUrea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834)Product Application DetailsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1 EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified
Product Description Description The prediluted antibody does not require any mixing, dilution, reconstitution, of titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 991 Gene Symbol CDC20 Species Human Immunogen Urea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834) Product Application Details Applications Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10 mM PBS with 0.05% BSA
DescriptionThe prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining.HostMouseGene ID991Gene SymbolCDC20SpeciesHumanImmunogenUrea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834)Product Application DetailsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1 EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Target Molecular Weight	55 kDa
titration; the antibody is ready-to-use and optimized for staining.HostMouseGene ID991Gene SymbolCDC20SpeciesHumanImmunogenUrea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834)Product Application DetailsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11 EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
Gene ID991Gene SymbolCDC20SpeciesHumanImmunogenUrea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834)Product Application DetailsImmunohistochemistry, Immunohistochemistry-ParaffinApplicationsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1r EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining.
Gene SymbolCDC20SpeciesHumanImmunogenUrea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834)Product Application DetailsApplicationsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1r EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Mouse
SpeciesHumanImmunogenUrea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834)Product Application DetailsApplicationsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1r EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	991
ImmunogenUrea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834)Product Application DetailsApplicationsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1r EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	CDC20
Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1r EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human
ApplicationsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1r EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	Urea-denatured His6 Cdc20 human recombinant protein (Uniprot: Q12834)
Recommended DilutionsImmunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1r EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Application NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staini of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1r EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1r EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml
Optimal dilution for a specific application should be determined.	Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

www.novusbio.com



Images

Immunohistochemistry-Paraffin: Cdc20 Antibody (AR12) - IHC-Prediluted [NBP2-44406] - Formalin-fixed, paraffin-embedded human Tonsil stained with Cdc20 Antibody (AR12).



Immunohistochemistry-Paraffin: Cdc20 Antibody (AR12) - IHC-Prediluted [NBP2-44406] - Formalin-fixed, paraffin-embedded human Gastric Carcinoma stained with Cdc20 Antibody (AR12).







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-44406

H00000991-Q01-10ug	Recombinant Human Cdc20 GST (N-Term) Protein
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-44406

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

