Product Datasheet

MCAM/CD146 Antibody (SPM620) NBP2-44510-0.1mg

Unit Size: 0.1 mg

Store at 4C.

www.novusbio.com

technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-44510

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-44510



NBP2-44510-0.1mg


MCAM/CD146 Antibody (SPM620)

Product Information Unit Size 0.1 mg Concentration 0.2 mg/ml Storage Store at 4C. Clonality Monoclonal Clonality Monoclonal Clone SPM620 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 30 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to a-80C. Host Mouse Gene ID 4162 Gene Symbol MCAM Species Human, Rat Reactivity Notes Rat reactivity reported in scientific literature (PMID: 31811403). Immunohistochemistry. Immunofustochemistry. Immunofustochemistry. Immunofustochemistry. Paratifin. Immunofustochemistry. Paratifin. Immunofustochemistry. Paratifin. Istuentific method protein (Uniprot: P43121) Product Application Details <t< th=""><th></th><th>,</th></t<>		,
Concentration0.2 mg/mlStorageStor at 4C.ClonalityMonoclonalCloneSPM620Preservative0.05% Sodium AzideIsotypeIgG1 KappaPurityProtein A or G purifiedBuffer10 mM PBS with 0.05% BSATarget Molecular Weight130 kDaProduct Description200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488)Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostMouseGene ID4162Gene SymbolMCAMSpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenRecombinant human MCAM/CD146 protein (Uniprot: P43121)Product Application DetailsFlow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunofluorescence 0.5 - 1.0 ug/mlApplication NotesFlow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunohistochemistry/Immunofluorescence, 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in TM EDTA, pH 8.0, of for 45 min at 95 C followed by cooling at RT for 20 minutes.	Product Information	
StorageStore at 4C.ClonalityMonoclonalCloneSPM620Preservative0.05% Sodium AzideIsotypeIgG1 KappaPurityProtein A or G purifiedBuffer10 mM PBS with 0.05% BSATarget Molecular Weight130 kDaProduct DescriptionDescriptionDescription200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (MBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostMouseGene ID4162Gene SymbolMCAMSpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenRecombinant human MCAM/CD146 protein (Uniprot: P43121)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunofluorescence -1 zu g/ml, Immunohistochemistry, Immunohistochemistry, Immunofluorescence -1 zu g/ml, Immunohistochemistry, Immunohistochemistry, I	Unit Size	0.1 mg
ClonalMonoclonalCloneSPM620Preservative0.05% Sodium AzideIsotypeIgG1 KappaPurityProtein A or G purifiedBuffer10 mM PBS with 0.05% BSATarget Molecular Weight130 kDaProduct Description200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (MBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostMouseGene ID4162Gene SymbolMCAMSpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403). ImmunogenRecombinant human MCAM/CD146 protein (Uniprot: P43121)Product Application DetailsApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunofluorescence 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of for 45 min at 95C followed by cooling at RT for 20 minutes.	Concentration	0.2 mg/ml
CloneSPM620Preservative0.05% Sodium AzideIsotypeIgG1 KappaPurityProtein A or G purifiedBuffer10 mM PBS with 0.05% BSATarget Molecular Weight130 kDaProduct Description200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488)Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostMouseGene ID4162Gene SymbolMCAMSpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenFlow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry. Immunohistochemistry/Immunofluorescence, Immunohistochemistry/Immunofluorescence, Immunohistochemistry/Immunofluorescence 0.5 - 1.0 ug/mlApplication NotesFlow Cytometry 1-2 ug/million cells, Immunohistochemistry.Application NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of forrmalin-fixed tissues requires heating iissue sections in 1mM EDTA, pH 8.0, for 45 min 41 sto 56 followed by cooling at RT for 20 minutes.	Storage	Store at 4C.
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 130 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4162 Gene Symbol MCAM Species Human, Rat Reactivity Notes Rat reactivity reported in scientific literature (PMID: 31811403). Immunogen Flow Cytometry . Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence Recommended Dilutions Flow Cytometry 1-2 ug/million cells, Immunohistochemistry. Parafin 1-2 ug/mi, Immunofluorescence 0.5 - 1.0 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating issue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clonality	Monoclonal
Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 130 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4162 Gene Symbol MCAM Species Human, Rat Reactivity Notes Rat reactivity reported in scientific literature (PMID: 31811403). Immunogen Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, Immunohistochemistry, Paraffin 1-2 ug/million cells, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Paraffin, Immunohistochemistry, Paraffin 1-2 ug/mil for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 mit at 95 followed by cooling at RT for 20 minutes.	Clone	SPM620
Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 130 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. -80C. Host Mouse Gene ID 4162 Gene Symbol MCAM Species Human, Rat Reactivity Notes Rat reactivity reported in scientific literature (PMID: 31811403). Immunogen Recombinant human MCAM/CD146 protein (Uniprot: P43121) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Paraffin 1-2 ug/ml, Inmunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml Application Notes Immunohistochemistry (Formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	0.05% Sodium Azide
Buffer10 mM PBS with 0.05% BSATarget Molecular Weight130 kDaProduct Description200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostMouseGene ID4162Gene SymbolMCAMSpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunofluorescence 1.2 ug/ml, ImmunofluorescenceRecommended DilutionsFlow Cytometry 1.2 ug/million cells, Immunohistochemistry, Immunofluorescence 1.2 ug/ml, Immunohistochemistry, Paraffin 1-2 ug/ml, Immunofluorescence 1.2 ug/ml, Immunohistochemistry, Paraffin 1-2 ug/ml, Immunofluorescence 1.2 ug/ml, Inmunohistochemistry, Paraffin 1-2 ug/ml, Immunofluorescence 1.2 ug/ml, Inmunohistochemistry, Para	Isotype	IgG1 Kappa
Target Molecular Weight130 kDaProduct Description200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostMouseGene ID4162Gene SymbolMCAMSpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenRecombinant human MCAM/CD146 protein (Uniprot: P43121)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Paraffin 1-2 ug/million cells, Immunohistochemistry, Par	Purity	Protein A or G purified
Product Description Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4162 Gene Symbol MCAM Species Human, Rat Reactivity Notes Rat reactivity reported in scientific literature (PMID: 31811403). Immunogen Recombinant human MCAM/CD146 protein (Uniprot: P43121) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Paraffin 1-2 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10 mM PBS with 0.05% BSA
Description200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostMouseGene ID4162Gene SymbolMCAMSpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/ml, Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/ml, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Paraffin 1-2 ug/ml, Immunofluorescence 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Target Molecular Weight	130 kDa
Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostMouseGene ID4162Gene SymbolMCAMSpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenRecombinant human MCAM/CD146 protein (Uniprot: P43121)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, Immunofluorescence 1-2 ug/ml, Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/ml, Immunofluorescence 0.5 - 1.0 ug/mlApplication NotesFlow Cytometry (Formalin-fixed tissues requires heating tissue sections in 1MM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
Gene ID4162Gene SymbolMCAMSpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenRecombinant human MCAM/CD146 protein (Uniprot: P43121)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, ImmunofluorescenceRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Inmunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-11488) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to
Gene SymbolMCAMSpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenRecombinant human MCAM/CD146 protein (Uniprot: P43121)Product Application DetailsApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, ImmunofluorescenceRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunohistochemistry, Immunofluorescence 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Mouse
SpeciesHuman, RatReactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenRecombinant human MCAM/CD146 protein (Uniprot: P43121)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, ImmunofluorescenceRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunohistochemistry	Gene ID	4162
Reactivity NotesRat reactivity reported in scientific literature (PMID: 31811403).ImmunogenRecombinant human MCAM/CD146 protein (Uniprot: P43121)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, ImmunofluorescenceRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunohistochemistry, Paraffin 1-2 ug/mi, Immunofluorescence 0.5 - 1.0 ug/miApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	MCAM
ImmunogenRecombinant human MCAM/CD146 protein (Uniprot: P43121)Product Application DetailsApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, ImmunofluorescenceRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/ml, Immunofluorescence 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human, Rat
Product Application Details Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence Recommended Dilutions Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Immunofluorescence 0.5 - 1.0 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Reactivity Notes	Rat reactivity reported in scientific literature (PMID: 31811403).
ApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, ImmunofluorescenceRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/ml, Immunofluorescence 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	Recombinant human MCAM/CD146 protein (Uniprot: P43121)
Immunohistochemistry, Immunohistochemistry-Paraffin, ImmunofluorescenceRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/ml, Immunofluorescence 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/ml, Immunofluorescence 0.5 - 1.0 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	
of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-
	Application Notes	of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 8.0, for 45 min at 95C followed by cooling at RT for 20 minutes.



Images

Immunohistochemistry-Paraffin: MCAM/CD146 Antibody (SPM620) [NBP2-44510] - Human Melanoma stained with MCAM Monoclonal Antibody (SPM620)



Publications

Pan J, Deng J, Yu L, et al. Investigating the repair of alveolar bone defects by gelatin methacrylate hydrogelsencapsulated human periodontal ligament stem cells J Mater Sci Mater Med 2019-12-05 [PMID: 31811403] (IF/IHC, Rat)





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-44510-0.1mg

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP1-88132PEP	MCAM/CD146 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-44510

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

