

Product Datasheet

Cyclin B1 Antibody (CCNB1/1098)

NBP2-44687-0.1mg

Unit Size: 0.1 mg

Store at 4C.

www.novusbio.com



technical@novusbio.com

Reviews: 1


Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-44687

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-44687




NBP2-44687-0.1mg

Cyclin B1 Antibody (CCNB1/1098)


Product Information	
Unit Size	0.1 mg
Concentration	0.2 mg/ml
Storage	Store at 4C.
Clonality	Monoclonal
Clone	CCNB1/1098
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS with 0.05% BSA
Product Description	
Description	200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-47893) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	891
Gene Symbol	CCNB1
Species	Human, Mouse
Specificity/Sensitivity	It recognizes a protein of 55-62kDa, identified as cyclin B1. In mammals, cyclin B associates with inactive p34cdc2, which facilitates phosphorylation of p34cdc2 at aa 14Thr and 15Tyr. This maintains the inactive state until the end of G2-phase. The inactive cyclin B-p34cdc2 complex continues to accumulate in the cytoplasm until the completion of DNA synthesis, when Cdc25, a specific protein phosphatase, dephosphorylates aa 14Thr and 15Tyr of p34cdc2 rendering the complex active at the G2/M boundary. This mitotic kinase complex remains active until the metaphase/anaphase transition when cyclin B is degraded. This degradation process is ubiquitin-dependent and is necessary for the cell to exit mitosis. So, cyclin B-p34cdc2 plays a critical role in G2 to M transition.
Immunogen	Recombinant human full-length Cyclin B1 protein (Uniprot: P14635)
Product Application Details	
Applications	Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1-2 ug/ml
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

Images


Immunocytochemistry/Immunofluorescence: Cyclin B1 Antibody (CCNB1/1098) [NBP2-44687] - MCF-7 cells stained with CF555 Cyclin B1 Antibody (CCNB1/1098) (nuclei/nucleoli, red) and CF488A phalloidin (actin filaments, green). DNA is stained with Hoechst (blue).




Immunohistochemistry-Paraffin: Cyclin B1 Antibody (CCNB1/1098) [NBP2-44687] - Human Tonsil stained with Cyclin B1 Monoclonal Antibody (CCNB1/1098)




Immunohistochemistry-Paraffin: Cyclin B1 Antibody (CCNB1/1098) [NBP2-44687] - Human Endometrial Carcinoma stained with Cyclin B1 Monoclonal Antibody (CCNB1/1098)




Immunohistochemistry-Paraffin: Cyclin B1 Antibody (CCNB1/1098) [NBP2-44687] - Human Ovarian Carcinoma stained with Cyclin B1 Monoclonal Antibody (CCNB1/1098)



Immunohistochemistry-Paraffin: Cyclin B1 Antibody (CCNB1/1098)
[NBP2-44687] - Human Prostate Carcinoma stained with Cyclin B1
Monoclonal Antibody (CCNB1/1098)



Immunohistochemistry-Paraffin: Cyclin B1 Antibody (CCNB1/1098)
[NBP2-44687] - Human Testicular Carcinoma stained with Cyclin B1
Monoclonal Antibody (CCNB1/1098)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-44687-0.1mg

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP1-98919-20ug	Recombinant Human Cyclin B1 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-44687

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

