## **Product Datasheet**

# CD79A Antibody (JCB117) - IHC-Prediluted NBP2-44736

Unit Size: 7 ml

Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-44736

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-44736




| CD79A Antibody (JCB117) - IHC-Prediluted |  |
|--|--|
| Product Information                      |  |
| Unit Size                                | 7 ml   |
| Concentration                            | Please see the vial label for concentration. If unlisted please contact technical services.  |
| Storage                                  | Store at 4C.   |
| Clonality                                | Monoclonal   |
| Clone                                    | JCB117   |
| Preservative                             | 0.05% Sodium Azide   |
| Isotype                                  | IgG1 Kappa   |
| Purity                                   | Protein A or G purified  |
| Buffer                                   | 10 mM PBS with 0.05% BSA   |
| Target Molecular Weight                  | 44 kDa   |
| Product Description                      |  |
| Description                              | The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining.  |
| Host                                     | Mouse  |
| Gene ID                                  | 973  |
| Gene Symbol                              | CD79A  |
| Species                                  | Human  |
| Marker                                   | B-Cell Marker  |
| Specificity/Sensitivity                  | A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. |
| Immunogen                                | A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein.  |
| Product Application Details              |  |
| Applications                             | Immunohistochemistry, Immunohistochemistry-Paraffin  |
| Recommended Dilutions                    | Immunohistochemistry, Immunohistochemistry-Paraffin 1:100-1:200  |
| Application Notes                        | Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.  |



### **Images**

Immunohistochemistry-Paraffin: CD79A Antibody (JCB117) - IHC-Prediluted [NBP2-44736] - Formalin-fixed, paraffin-embedded human Tonsil stained with CD79A Antibody (JCB117).





#### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

#### **Products Related to NBP2-44736**

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

NBP2-60209-50ug Recombinant Human CD79A His Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-44736

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

