## **Product Datasheet**

### CD8 Antibody (C8/468) - IHC-Prediluted NBP2-45323

Unit Size: 7 ml

Store at 4C.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-45323

Updated 10/23/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-45323



#### NBP2-45323


CD8 Antibody (C8/468) - IHC-Prediluted

Product Information	
Unit Size	7 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C.
Clonality	Monoclonal
Clone	C8/468
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS with 0.05% BSA
Target Molecular Weight	32 kDa
Product Description	
Description	The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining.
Host	Mouse
Gene ID	925
Gene Symbol	CD8A
Species	Human
Marker	Cytotoxic / Suppressor T-Cell Marker
Specificity/Sensitivity	CD8 Antibody (C8/468) - IHC-Prediluted recognizes CD8, a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha/beta) or as a homodimer (CD8 alpha/alpha). A majority of thymocytes and a subpopulation of mature T cells and NK cells express CD8a. CD8 binds to MHC class 1 and through its association with protein tyrosine kinase p56lck plays a role in T cell development and activation of mature T cells. For mature T-cells, CD4 and CD8 are mutually exclusive, so anti-CD8 is generally used in conjunction with anti-CD4. It is a useful marker for distinguishing helper/inducer T-lymphocytes, and most peripheral T-cell lymphomas are CD4+/CD8 Anaplastic large cell lymphoma is usually CD4+ and CD8-, and in T- lymphoblastic lymphoma/leukemia, CD4 and CD8 are often co-expressed. CD8 is also found in littoral cell angioma of the spleen.
Immunogen	Human CD8 recombinant protein (Uniprot: P01732)
Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
<b>Recommended Dilutions</b>	Immunohistochemistry, Immunohistochemistry-Paraffin 1:100 - 1:200
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.



#### Images

Immunohistochemistry-Paraffin: CD8 Antibody (C8/468) - IHC-Prediluted [NBP2-45323] - Formalin-fixed, paraffin-embedded human lymph node stained with CD8 Mouse Monoclonal Antibody (C8/468).



Immunohistochemistry-Paraffin: CD8 Antibody (C8/468) - IHC-Prediluted [NBP2-45323] - Formalin-fixed, paraffin-embedded human tonsil stained with CD8 Mouse Monoclonal Antibody (C8/468).





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-45323

NBP2-34039PEP	CD8 Recombinant Protein Antigen
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-45323

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

