

Product Datasheet

VEGF-C Antibody (OTI4A1)

NBP2-45404

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-45404

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-45404



NBP2-45404

VEGF-C Antibody (OTI4A1)


Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	OTI4A1
Preservative	0.02% Sodium Azide
Isotype	IgG1
Purity	Immunogen affinity purified
Buffer	PBS (pH 7.3), 1.0% BSA and 50% Glycerol
Target Molecular Weight	46.7 kDa

Product Description	
Host	Mouse
Gene ID	7424
Gene Symbol	VEGFC
Species	Human, Mouse, Rat
Reactivity Notes	Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.
Immunogen	Human recombinant protein fragment corresponding to amino acids 112-227 of human VEGFC (NP_005420) produced in E.coli.


Product Application Details	
Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1:200, Immunohistochemistry, Immunohistochemistry-Paraffin

Images


Western Blot: VEGF-C Antibody (OTI4A1) [NBP2-45404] - Analysis of HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY VEGF-C.




Immunohistochemistry-Paraffin: VEGF-C Antibody (OTI4A1) [NBP2-45404] - Staining of paraffin-embedded Human prostate tissue within the normal limits using anti-VEGFC mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120C for 3min) (1:150)




Immunohistochemistry-Paraffin: VEGF-C Antibody (OTI4A1) [NBP2-45404] - Staining of paraffin-embedded Carcinoma of Human liver tissue using anti-VEGFC mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120C for 3min) (1:150)




Immunohistochemistry-Paraffin: VEGF-C Antibody (OTI4A1) [NBP2-45404] - Staining of paraffin-embedded Carcinoma of Human pancreas tissue using anti-VEGFC mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120C for 3min) (1:150)




Immunohistochemistry-Paraffin: VEGF-C Antibody (OTI4A1) [NBP2-45404] - Staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-VEGFC mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120C for 3min) (1:150)




Immunohistochemistry-Paraffin: VEGF-C Antibody (OTI4A1) [NBP2-45404] - Staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-VEGFC mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120C for 3min) (1:150)




Immunohistochemistry-Paraffin: VEGF-C Antibody (OTI4A1) [NBP2-45404] - Staining of paraffin-embedded Human liver tissue within the normal limits using anti-VEGFC mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120C for 3min) (1:150)



Immunohistochemistry-Paraffin: VEGF-C Antibody (OTI4A1) [NBP2-45404] - Staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-VEGFC mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120C for 3min) (1:150)



Immunohistochemistry-Paraffin: VEGF-C Antibody (OTI4A1) [NBP2-45404] - Staining of paraffin-embedded Human tonsil within the normal limits using anti-VEGFC mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120C for 3min) (1:150)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-45404

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
DVE00	VEGF [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-45404

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

