Product Datasheet

Human Prostate Tissue Lysate (Adult Normal) NBP2-47059

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47059

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47059



NBP2-47059

Human Prostate Tissue Lysate (Adult Normal)

, (
Product Information	
Unit Size	0.1 mg
Concentration	Please contact technical services for concentration.
Storage	Store at -80C. Avoid freeze-thaw cycles.
Product Description	
Description	Normal Tissue Lysate
	Diagnosis: Normal prostate
	Sex: Male
	Age: 37
	Grade: N/A
	Stage: N/A
	Normal Tissue Pathology Data
	Location: Prostate
	Gross findings: No data available
	Microscopic Findings
	No data available
	Preparation Method
	Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4
Species	Human
Notes	The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately
Lysate Type	Tissue
Lysate Tissue	Prostate
Lysate Tissue Condition	Normal
Lysate Life Stage	Adult
Product Application Details	· · · · · · · · · · · · · · · · · · ·

Product Application Details



Application Notes	 These lysates have not been subjected to denaturing or reducing conditions. This allows the tissue or cell lysate to be used in a variety of applications; to study protein-protein interaction, ligand binding, ELISA, immunoprecipitation, 1D and 2D gel electrophoresis, and Western blotting for the detection of specific protein targets. For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel loading buffer with reducing agents may be required. Buffer requirements for performing protein-protein interaction and ligand binding studies can vary significantly from RIPA buffer and may require modifications. In most cases, tissue lysates in RIPA buffer can be used, directly in standard ELISA and immunoprecipitation assays. These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47059

NBP2-30169

Human Prostate Tissue MicroArray (Cancer)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47059

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

