Product Datasheet

Human Prostate Tissue Lysate (Adult Normal) NBP2-47063

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47063

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47063



NBP2-47063

Human Prostate Tissue Lysate (Adult Normal)

Human Prostate Tissue Lysate (Adult Normal)	
0.1 mg	
Please contact technical services for concentration.	
Store at -80C. Avoid freeze-thaw cycles.	
Normal Tissue Lysate	
Diagnosis: Normal prostate	
Sex: Male	
Age: 32	
Grade: N/A	
Stage: N/A	
Normal Tissue Pathology Data	
Location: Prostate	
Gross findings: No data available	
Microscopic Findings	
No data available	
Preparation Method	
Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4	
Human	
The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately	
Tissue	
Prostate	
Normal	
Adult	

Product Application Details



Application Notes

These lysates have not been subjected to denaturing or reducing conditions. This allows the tissue or cell lysate to be used in a variety of applications; to study protein-protein interaction, ligand binding, ELISA, immunoprecipitation, 1D and 2D gel electrophoresis, and Western blotting for the detection of specific protein targets.

For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel loading buffer with reducing agents may be required.

Buffer requirements for performing protein-protein interaction and ligand binding studies can vary significantly from RIPA buffer and may require modifications. In most cases, tissue lysates in RIPA buffer can be used, directly in standard ELISA and immunoprecipitation assays.

These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-47063

NBP2-30169

Human Prostate Tissue MicroArray (Cancer)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47063

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

