Product Datasheet

Stomach Matched Tumor and Normal Tissue Lysate NBP2-47082

Unit Size: 2 Vials

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com

technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47082

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47082



NBP2-47082

Stomach Matched Tumor and Normal Tissue Lysate

Product Information Unit Size 2 Vials Concentration mg/ml Storage Store at -80C. Avoid freeze-thaw cycles. Product Description 1 x Human stomach tumor tissue lysate (1 mg/ml, 100 ug/vial) 1 x Human stomach normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Matched Tumor & Normal Tissue Lysate Set Diagnosis: Adenocarcinoma Sex: Male Sex: Male	g/vial)
Concentration mg/ml Storage Store at -80C. Avoid freeze-thaw cycles. Product Description 1 x Human stomach tumor tissue lysate (1 mg/ml, 100 ug/vial) 1 x Human stomach normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) 1 x Human stomach normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Diagnosis: Adenocarcinoma	g/vial)
Storage Store at -80C. Avoid freeze-thaw cycles. Product Description 1 x Human stomach tumor tissue lysate (1 mg/ml, 100 ug/vial) 1 x Human stomach normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) 1 x Human stomach normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Diagnosis: Adenocarcinoma	g/vial)
Product Description Description 1 x Human stomach tumor tissue lysate (1 mg/ml, 100 ug/vial) 1 x Human stomach normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) 1 x Human stomach normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Diagnosis: Adenocarcinoma	g/vial)
Description 1 x Human stomach tumor tissue lysate (1 mg/ml, 100 ug/vial) 1 x Human stomach normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Matched Tumor & Normal Tissue Lysate Set Diagnosis: Adenocarcinoma	g/vial)
1 x Human stomach normal tissue lysate (matched) (1 mg/ml, 100 u Matched Tumor & Normal Tissue Lysate Set Diagnosis: Adenocarcinoma	g/vial)
Diagnosis: Adenocarcinoma	
Sex: Male	
Age: 48	
Grade: N/A	
Stage: IIIA, T3N1M0	
Tumor Pathology Data	
Location: Gastric antrum	
Gross findings: Tumor located at the pyloric antrum. Tumor measure cm x 4 cm in diameter. Lesion is large, has an irregular profile, show and interrupts the gastric folds. Paragastric lymph nodes: 6.	
Microscopic Findings	
The tumor is composed of nests of epithelial cells forming the gland, chromatin, and nuclear pleomorphism. The tumor cells also are arrant trebecular patterns or cords, and invade the muscular wall and seroes necrosis is present, as well as lymphocytic response. Lymph nodes a positive.	nged in sa. Tumor
Preparation Method	
Tissue specimens are homogenized in modified RIPA buffer to obtai proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4	
Species Human	
Notes The vial is provided with a 10% overfill. Maximum recovery can be of centrifuging the vial briefly to collect any solution on the cap and tube	
This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivati treated as a potentially infectious agent and disposed of appropriate	ves must be



	5 1	
Lysate Type	Matched Tumor and Normal Tissue	
Lysate Tissue	Stomach	
Lysate Life Stage	Adult	
Product Application Details		
Application Notes	These lysates have not been subjected to denaturing or reducing conditions. This allows the tissue or cell lysate to be used in a variety of applications; to study protein-protein interaction, ligand binding, ELISA, immunoprecipitation, 1D and 2D gel electrophoresis, and Western blotting for the detection of specific protein targets. For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel loading buffer with reducing agents may be required. Buffer requirements for performing protein-protein interaction and ligand binding studies can vary significantly from RIPA buffer and may require modifications. In most cases, tissue lysates in RIPA buffer can be used, directly in standard ELISA and immunoprecipitation assays. These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use.	

Publications

Ara H, Subedi U, Sharma P et al. Alteration of Cellular Energy Metabolism through LPAR2-Axin2 Axis in Gastric Cancer Biomolecules 2022-12-02 [PMID: 36551233]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47082

NBP2-30308

Human Stomach Tissue MicroArray (Cancer)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47082

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

