Product Datasheet

Liver Matched Tumor and Normal Tissue Lysate NBP2-47087-2Vial

Unit Size: 2 Vial

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47087

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47087



NBP2-47087-2Vial

Liver Matched Tumor and Normal Tissue Lysate

Unit Size 2 Vial Concentration mg/ml Storage Store at -80C. Avoid freeze-thaw cycles. Product Description 1 x Human Liver tumor tissue lysate (1 mg/ml, 100 ug/vial) 1 x Human Liver normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) 1 x Human Liver normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) 1 x Human Liver normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) 1 x Human Liver normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Pactor Natched Tumor & Normal Tissue Lysate Set Diagnosis: Hepatocellular carcinoma Sex: Male Age: 55 Grade: 2 Grade: 2 Stage: N/A Tumor Pathology Data Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic.ccipilaries and infiltration of lymphocytes. Necroses and blod vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, PI 7.4: 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer. 1 mM EDTA; 1 ug/ml Pepattin-A; 0.1% SDS; 0.25%, Na	Product Information	
Storage Store at -80C. Avoid freeze-thaw cycles. Product Description I x Human Liver tumor tissue lysate (1 mg/ml, 100 ug/vial) 1 x Human Liver normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Description 1 x Human Liver normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Matched Tumor & Normal Tissue Lysate Set Diagnosis: Hepatocellular carcinoma Sex: Male Age: 55 Grade: 2 Grade: 2 Stage: N/A Tumor Pathology Data Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surronding stroma show shyperplastic capilalries and infiltration of ymphocytes. Necroses and blood vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, PT.4; 1 ug/ml Apottinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 md MSA; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 md MSA; 0.1% JDS; 0.25% Na deoxycholate; 1 ug/ml PRECAUTIONS when handing. Human tissue derivatives must be treated as a potentially intectious agenit and disposed of appropritaley	Unit Size	2 Vial
Product Description Description 1 x Human Liver tumor tissue lysate (1 mg/ml, 100 ug/vial) 1 x Human Liver normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Matched Tumor & Normal Tissue Lysate Set Diagnosis: Hepatocellular carcinoma Sex: Male Age: 55 Grade: 2 Stage: N/A Tumor Pathology Data Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoil and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centifuged to clarify. Extraction 1: PBS, PI 7 4: 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstain-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM NaSVo4 Species Human Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifying the vial briefly to collect any solution on the cap and tube sides.		
Description 1 x Human Liver tumor tissue lysate (1 mg/ml, 100 ug/vial) 1 x Human Liver normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Matched Tumor & Normal Tissue Lysate Set Diagnosis: Hepatocellular carcinoma Sex: Male Age: 55 Grade: 2 Stage: N/A Tumor Pathology Data Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins. and centrifuged to clarify. Extraction 1: PBS, PI 7.4, 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM NaSVO4 Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. <th>Storage</th> <th>Store at -80C. Avoid freeze-thaw cycles.</th>	Storage	Store at -80C. Avoid freeze-thaw cycles.
1 x Human Liver normal tissue lysate (matched) (1 mg/ml, 100 ug/vial) Matched Tumor & Normal Tissue Lysate Set Diagnosis: Hepatocellular carcinoma Sex: Male Age: 55 Grade: 2 Stage: N/A Tumor Pathology Data Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoii and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Laupeptin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Papstain-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Laupeptin; 1 mM PMSF; 1 mM Na3VO4 Species Human Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HV 1/2, and HCV. Use UNVERSAAL PRECAUTIONS when handling. Human tissue derivatives	Product Description	
Diagnosis: Hepatocellular carcinoma Sex: Male Age: 55 Grade: 2 Stage: N/A Tumor Pathology Data Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tussue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoi and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and biod vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4 Species Human Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial biefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handing, Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately	Description	
Sex: Male Age: 55 Grade: 2 Stage: N/A Tumor Pathology Data Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4 Species Human Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately		Matched Tumor & Normal Tissue Lysate Set
Age: 55 Grade: 2 Stage: N/A Tumor Pathology Data Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleil and abundant. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, PH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4 Species Human Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately		Diagnosis: Hepatocellular carcinoma
Grade: 2 Stage: N/A Tumor Pathology Data Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4 Species Human Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS When handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately		Sex: Male
Stage: N/A Tumor Pathology Data Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4 Species Human Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately		Age: 55
Species Human Notes Human Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides.		Grade: 2
Location: Liver Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color. Microscopic findings Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed. Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4 Species Human Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately		Stage: N/A
Gross findings: Tumor is 10 cm in diameter, well demarcated. Cut section is soft, hemorrhagic, and gray-white in color.Microscopic findingsTissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma 		Tumor Pathology Data
hemorrhagic, and gray-white in color.Microscopic findingsTissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed.Preparation MethodTissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify.Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4SpeciesHumanNotesThe vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately		Location: Liver
Tissue sections show proliferation of malignant epithelial cell clusters with round nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed.Preparation MethodTissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4SpeciesHumanNotesThe vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately		
nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and blood vessel invasion not revealed.Preparation MethodTissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4SpeciesHumanNotesThe vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately		Microscopic findings
SpeciesNotesHumanNotesThis material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately		nuclei, predominant nucleoli and abundant cytoplasm. Nuclear chromatin is coarse and irregularly distributed. Mitoses are evident. The surrounding stroma shows hyperplastic capillaries and infiltration of lymphocytes. Necroses and
proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na deoxycholate; 1 ug/ml Leupeptin; 1 mM PMSF; 1 mM Na3VO4SpeciesHumanNotesThe vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be 		Preparation Method
Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately		proteins, and centrifuged to clarify. Extraction 1: PBS, pH 7.4; 1 ug/ml Aprotinin; 1 mM NaF Modified RIPA Buffer: 1 mM EDTA; 1 ug/ml Pepstatin-A; 0.1% SDS; 0.25% Na
centrifuging the vial briefly to collect any solution on the cap and tube sides. This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately	Species	Human
UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately	Notes	centrifuging the vial briefly to collect any solution on the cap and tube sides.
Lysate Type Matched Tumor and Normal Tissue		UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately
	Lysate Type	Matched Tumor and Normal Tissue



Lysate Tissue	Liver
Lysate Life Stage	Adult
Product Application Details	
Application Notes	These lysates have not been subjected to denaturing or reducing conditions. This allows the tissue or cell lysate to be used in a variety of applications; to study protein-protein interaction, ligand binding, ELISA, immunoprecipitation, 1D and 2D gel electrophoresis, and Western blotting for the detection of specific protein targets. For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel loading buffer with reducing agents may be required. Buffer requirements for performing protein-protein interaction and ligand binding studies can vary significantly from RIPA buffer and may require modifications. In most cases, tissue lysates in RIPA buffer can be used, directly in standard ELISA and immunoprecipitation assays. These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47087-2Vial

NBP2-30221

Human Liver Tissue MicroArray (Cancer metastasis)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47087

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

