Product Datasheet

Human IgG Fc Isotype Control NBP2-47132

Unit Size: 1 ml

Store at 4C.

www.novusbio.com

technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47132

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47132



NBP2-47132

Human IgG Fc Isotype Control

| Product Information | |
|------------------------------|---|
| Unit Size | 1 ml |
| Concentration | 1 mg/ml |
| Storage | Store at 4C. |
| Clonality | Polyclonal |
| Preservative | 0.09% Sodium Azide |
| Isotype | IgG |
| Purity | Affinity purified |
| Buffer | Phosphate Buffered Saline |
| Product Description | |
| Description | A human expressed protein corresponding to IgGFc. |
| Host | Human |
| Species | Human |
| Preparation Method | Purified normal human IgG is isolated using fractionation and DEAE chromatography. Fc fragments were produced by papain digestion and affinity chromatography. Protein concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis the IgG was shown to 1) react with antiserum specific for human IgGFc, 2) not react with antiserum specific for IgA, IgM, kappa or lambda chain and 3) produce a single precipitin arc with antiserum against human serum identical to that produced with antiIgG-Fc. |
| Notes | Protein concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis the IgG was shown to 1) react with antiserum specific for human IgGFc, 2) not react with antiserum specific for IgA, IgM, kappa or lambda chain and 3) produce a single precipitin arc with antiserum against human serum identical to that produced with antilgG-Fc. |
| Product Application Details | |
| Applications | ELISA, Immunoprecipitation |
| Recommended Dilutions | ELISA, Immunoprecipitation |
| Application Notes | Serum or plasma from donor has been tested and found negative for HIV-1 antigen, antibodies for HIV and HCV, and non-reactive for HBsAg, HIV-1 RNA, and HCV RNA. Not tested for AntiHBc. HANDLE AS IF CAPABLE OF TRANSMITTING INFECTIOUS AGENTS. |

Publications

Vogl T, Klompus S, Leviatan S Et Al. Population-wide diversity and stability of serum antibody epitope repertoires against human microbiota Nature medicine 2021-07-19 [PMID: 34282338]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47132

NB7446

Goat anti-Human IgG Fc Secondary Antibody

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Isotype Controls are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47132

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

