Product Datasheet

HLA-DR Antibody (LN-3) [Alexa Fluor® 405] NBP2-47670AF405

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47670AF405

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47670AF405



NBP2-47670AF405

HLA-DR Antibody (LN-3) [Alexa Fluor® 405]

| Product Information Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone LN-3 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Host Mouse Gene Bymbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody recognizes a different epitope than the LN3 monoclonal antibody recognizes a different epitope than the LN3 monoclonal antibody. and these antibodies do not cross-block binding to each other coefficience and coefficience philos and coefficience and coefficience and coefficience and coefficience philos and coefficience philos and coefficience and coefficience philos and coefficience philos and coefficience and coefficience philos and coefficience and coefficience philos and coefficience philos and coefficience philos and coefficience | | · · · · · · · · · · · · · · · · · · · |
|---|-----------------------------|---|
| Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone LN-3 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L24 antibody recognizes a different epitope than. It is an onclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a neterodimeric cell surface glycorotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dentific cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for eflicient petide presentation to CD4+T cells. It is an excellent histocytic marker in paraffin sections producing intense staining. True histocyt | Product Information | |
| services. Storage Store at 4C in the dark. Clonality Monoclonal Clone LN-3 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer Sodium Borate Product Description Mouse Gene ID 3122 Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DR1 antigen, a member of MHC class II molecules. It does not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophagex and CD4 molecules. HLA-DR is a critical for efficient pepide presentation to CD4+T comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophagex and CD4 molecules, HLA-DR is critical for efficient pepide presentation to CD4+T comprise of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, sotivated T-cells, monocytes/macrophagex and CD4 molecules, HLA-DR is critical for efficient pepide presentation to CD4+T corour on avarie/of epithelial cells and their corresponding | Unit Size | 0.1 ml |
| Clonality Monoclonal Clone LN-3 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 3122 Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross-tock binding to each orbit res respective epitopes. HLA-DR is a heterodimeric cell surface glycorptein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histicocytic neoplasms are similarly positive. HLA-DR antigens also court on a variety of epithelial cells and their corresponding neoplastic court on a variety of epithelial cells and their corresponding neoplastic court or a variety of epithelial cells and their corresponding neoplastic court or a variety of epithelial cells and their corresponding neoplastic court or a variety of epithelial cells and their corresponding neoplastic court or avariety of epithel | Concentration | • |
| Clone LN-3 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description House Gene ID 3122 Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dentitic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules. HLA-DE is critical for efficient peptide presentation to CD4 + Cells Tiles expression is related to tumor microenvironment and predict adverse outcome in diffuse large B-cell lymphoma. Immunogen Activated human peripheral blood mononuclear cells Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunofluores | Storage | Store at 4C in the dark. |
| Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 3122 Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross-relow thin HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR his a heterodimeric cells surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histocytic neoplasms are similarly positive. HLA-DR is critical for efficient peptide presentation to CD4+ T cells. To cass of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma. Immunogen Activated human peripheral blood mononuclear cells Product Application Details Western Blot, Fl | Clonality | Monoclonal |
| IsotypeIgG2b KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID3122Gene SymbolHLA-DRASpeciesHuman, Monkey, Mouse (Negative)Reactivity NotesDoes not react with Mouse.Specificity/SensitivityThis monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recorgizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimetric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta [iight) chain. It is expressed on B-cells, activated T-cells. monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic marker in parafin sections producing intense staining. True histiccytic marker in parafin sections producing intense staining. The histiccytic moles and see allymphoma.ImmunogenActivated human peripheral blood mononuclear cellsProduct Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry/Immunofluorescence, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready | Clone | LN-3 |
| Conjugate Alexa Fluor 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 3122 Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta [light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4 T counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma. Immunogen Activated human peripheral blood mononuclear cells Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, CyTOF-ready | Preservative | 0.05% Sodium Azide |
| Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 3122 Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is a netwoellent histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts. Loss of HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent HLA-DR is a represion is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma. Immunogen Activated human peripheral blood mononuclear cells Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry. Immunohistochemistry/ Immunofluorescence, CyTOF-ready. | Isotype | IgG2b Kappa |
| Buffer 50mM Sodium Borate Product Description Mouse Gene ID 3122 Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DP. The L243 antibody recognizes a different peitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma. Immunogen Xetivated human peripheral blood mononuclear cells Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry./Immunofluorescence, Immunohistochemistry. Immunofluorescence, CyTOF-ready. Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry.Paraffin, Immunofluorescence, CyTOF-ready. | Conjugate | Alexa Fluor 405 |
| Product Description Host Mouse Gene ID 3122 Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4 + T cells. It is an excellent histiocytic marker in paraffin sections producing intense staining. True histiocytic moleals and their corresponding neoplastic counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma. Immunogen Activated human peripheral blood mononuclear cells Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, CyTOF-ready. | Purity | Protein A or G purified |
| HostMouseGene ID3122Gene SymbolHLA-DRASpeciesHuman, Monkey, Mouse (Negative)Reactivity NotesDoes not react with Mouse.Specificity/SensitivityThis monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies to not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic marker in paraffin sections producing intense staining. True histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma.ImmunogenActivated human peripheral blood mononuclear cellsProduct Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence | Buffer | 50mM Sodium Borate |
| Gene ID 3122 Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiccytic marker in paraffin sections producing intense staining. True histiccytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma. Immunogen Activated human peripheral blood mononuclear cells Application Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-ready. Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Paraffin, Immunofluorescence, CyTOF-ready | Product Description | |
| Gene Symbol HLA-DRA Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic marker in paraffin sections producing intense staining. True histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma. Immunogen Activated human peripheral blood mononuclear cells Product Application Details Western Blot, Flow Cytometry, Immunofucrescence, Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, CyTOF-ready Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunofluorescence, CyTOF-ready | Host | Mouse |
| Species Human, Monkey, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic marker in paraffin sections producing intense staining. True histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma. Immunogen Activated human peripheral blood mononuclear cells Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry/ Immunofluorescence, Immunohistochenistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready | Gene ID | 3122 |
| Reactivity NotesDoes not react with Mouse.Specificity/SensitivityThis monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma.ImmunogenActivated human peripheral blood mononuclear cellsProduct Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry. Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready | Gene Symbol | HLA-DRA |
| Specificity/SensitivityThis monoclonal antibody reacts with a 28kDa chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma.ImmunogenActivated human peripheral blood mononuclear cellsProduct Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunofluorescence, CyTOF-ready | Species | Human, Monkey, Mouse (Negative) |
| member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic | Reactivity Notes | Does not react with Mouse. |
| Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready | Specificity/Sensitivity | member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the LN3 monoclonal antibody, and these antibodies do not cross-block binding to each others respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic marker in paraffin sections producing intense staining. True histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma. |
| ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready | Immunogen | Activated human peripheral blood mononuclear cells |
| Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready | Product Application Details | |
| Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready | Applications | Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, |
| Application NotesOptimal dilution of this antibody should be experimentally determined. | Recommended Dilutions | Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, |
| | Application Notes | Optimal dilution of this antibody should be experimentally determined. |



Images

HLA-DR Antibody (LN-3) [Alexa Fluor® 405] [NBP2-47670AF405] - Vial of Alexa Fluor 405 conjugated antibody. Alexa Fluor 405 is optimally excited at 401 nm by the Violet laser (405 nm) and has an emission maximum of 421 nm.



www.novusbio.com





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47670AF405

NBP1-43317AF405Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [Alexa Fluor®
405]210-TA-005TNF-alpha [Unconjugated]M6000B-1IL-6 [HRP]285-IF-100IFN-gamma [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47670AF405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

