Product Datasheet

TNF-alpha Antibody (J2D10) [DyLight 650] NBP2-47677C

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47677C

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-47677C



NBP2-47677C

TNF-alpha Antibody (J2D10) [DyLight 650]

Unit Size	Trui -aipria Aritibody (32010) [Dyi	Tri -alpha Ahlibody (02D10) [DyEight 050]	
Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark.	Product Information		
Storage Store at 4C in the dark. Clonality Monoclonal Clone JZD10 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 7124 Gene Symbol TNF Species Human, Rat, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This antibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (Impnhotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells. Thr-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Notes Product Application Details Applications	Unit Size	0.1 ml	
Clone J2D10 Preservative 0.05% Sodium Azide Isotype 1gG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 7124 Gene Symbol TNF Species Human, Rat, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This antibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lipopolysacharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity, although it has little effect on many cultured normal human cells. ThR-alpha in appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophis and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Concentration	· ·	
Clone J2D10 Preservative 0.05% Sodium Azide Isotype 1gG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 7124 Gene Symbol TNF Species Human, Rat, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This antibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNFa.) alpha is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogachis chock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17Ka band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha acuses cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells. TNF-alpha appears to be directly toxic to vascular endo	Storage	Store at 4C in the dark.	
Preservative Isotype IgG1 Kappa	Clonality	Monoclonal	
IgG1 Kappa	Clone	J2D10	
Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 7124 Gene Symbol TNF Species Human, Rat, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This antibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lippolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injuly associated with endotoxemia. TNF-alpha iststs as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity, Although it has little effect on many cultured normal human cells, TNF-alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha inclosed stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Preservative	0.05% Sodium Azide	
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 7124 Gene Symbol TNF Species Human, Rat, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This ambibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF-alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Isotype	IgG1 Kappa	
Product Description	Conjugate	DyLight 650	
Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 7124 Gene Symbol TNF Species Human, Rat, Mouse (Negative) Does not react with Mouse. Specificity/Sensitivity This antibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lippoplysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF-alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophilis and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Purity	Protein A or G purified	
This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 7124 Gene Symbol TNF Species Human, Rat, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This antibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF-alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Buffer	50mM Sodium Borate	
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 7124 Gene Symbol TNF Species Human, Rat, Mouse (Negative) Does not react with Mouse. Specificity/Sensitivity This antibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF-alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophilis and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Product Description		
Gene Symbol TNF Species Human, Rat, Mouse (Negative) Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This antibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lipopolysaccharidestimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF-alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Description	volume of this product. The volume will be greater than or equal to the unit size	
Species	Host	Mouse	
Species	Gene ID	7124	
Reactivity Notes Does not react with Mouse. Specificity/Sensitivity This antibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF-alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Gene Symbol	TNF	
This antibody neutralizes Hu rTNFa mediated cytotoxicity of L929 cells and inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF-alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Species	Human, Rat, Mouse (Negative)	
inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF-alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. Immunogen Recombinant human TNF-alpha (Uniprot: P01375) Notes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Reactivity Notes	Does not react with Mouse.	
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Specificity/Sensitivity	inhibits tumor growth in mice. It protects mice against toxicity of HurTNFa. Tumor Necrosis Factor Alpha (TNF-alpha) is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF-alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF-alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF-alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF-alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF-alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF-alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of	
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Immunogen	Recombinant human TNF-alpha (Uniprot: P01375)	
Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	
	Product Application Details		
	• •		



	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-47677C

NBP1-43319C Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 650]

NBP2-35076-10ug Recombinant Human TNF-alpha Protein

M6000B-1 IL-6 [HRP]

210-TA-005 TNF-alpha [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47677C

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

