Product Datasheet

alpha-Smooth Muscle Actin Antibody (1A4 + ACTA2/791) [Alexa Fluor® 405] NBP2-47699AF405

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47699AF405

Updated 9/11/2024 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47699AF405

NBP2-47699AF405

alpha-Smooth Muscle Actin Antibody (1A4 + ACTA2/791) [Alexa Fluor® 405]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClone1A4 + ACTA2/791Preservative0.05% Sodium AzideIsotypeIgG2a Kappa/IgG2a KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionS9Gene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscle soft and myogenin are positive culd be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and anti-smooth muscle and anti-smooth muscle actin and are negative with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and anti-smooth muscle actin and are negative with anti-myogenin.ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/7791) (Uniprot: P62736)		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClone1A4 + ACTA2/791Preservative0.05% Sodium AzideIsotypeIgG2a Kappa/IgG2a KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description59Gene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin and sheletal muscle specific actin and myogenin and ant-smooth muscle actin and are negative with anti-muscle specific actin and myogenin (ACTA2/791) (Uniprot:ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalClone1A4 + ACTA2/791Preservative0.05% Sodium AzideIsotypeIgG2a Kappa/IgG2a KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description59HostMouseGene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle, however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific catin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific catin and anti-smooth muscle actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Unit Size	0.1 ml
ClonalMonoclonalClone1A4 + ACTA2/791Preservative0.05% Sodium AzideIsotypeIgG2a Kappa/IgG2a KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle, however, it does stain myofibroblasts and myoepitheilal cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of thabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcoma sare positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative vith anti-muscle specific actin and anti-smooth muscle actin and are negative vith anti-muscle specific actin and myogenin are positive. Leiomyosarcoma sare positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative vith anti-muscle specific actin and anti-smooth muscle actin and are negative vith anti-muscle specific actin and anti-smooth muscle actin and are negative vith anti-muscle specific actin and anti-smooth muscle actin and are negative vith anti-	Concentration	
Clone1A4 + ACTA2/791Preservative0.05% Sodium AzideIsotypeIgG2a Kappa/IgG2a KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cariac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG2a Kappa/IgG2a KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not scale and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and myoepithelial cells and are negative with anti-myogenin.ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/T91) (Uniprot:	Clonality	Monoclonal
IsotypeIgG2a Kappa/IgG2a KappaConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and <b< th=""><th>Clone</th><th>1A4 + ACTA2/791</th></b<>	Clone	1A4 + ACTA2/791
ConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and anti-smooth muscle actin and aconjugated to KLH (1	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Host Mouse Gene ID 59 Gene Symbol ACTA2 Species Human, Rat Reactivity Notes Predicted to show a broad reactivity. Marker Leiomyosarcoma Marker Specificity/Sensitivity Actin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-muscle specific actin and hyogenin. Immunogen N-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (144); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Isotype	IgG2a Kappa/IgG2a Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-myogenin.ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Conjugate	Alexa Fluor 405
Product Description Mouse Host Mouse Gene ID 59 Gene Symbol ACTA2 Species Human, Rat Reactivity Notes Predicted to show a broad reactivity. Marker Leiomyosarcoma Marker Specificity/Sensitivity Actin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-myogenin. Immunogen N-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Purity	Protein A or G purified
HostMouseGene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle 	Buffer	50mM Sodium Borate
Gene ID59Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-myogenin.ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Product Description	
Gene SymbolACTA2SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-myogenin.ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Host	Mouse
SpeciesHuman, RatReactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-myogenin.ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Gene ID	59
Reactivity NotesPredicted to show a broad reactivity.MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-myogenin.ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Gene Symbol	ACTA2
MarkerLeiomyosarcoma MarkerSpecificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-myogenin.ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Species	Human, Rat
Specificity/SensitivityActin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-myogenin.ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Reactivity Notes	Predicted to show a broad reactivity.
It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-myogenin.ImmunogenN-Terminal decapeptide of alpha-Smooth Muscle Actin and conjugated to KLH (1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Marker	Leiomyosarcoma Marker
(1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:	Specificity/Sensitivity	It is highly specific to actin from smooth muscles. This monoclonal antibody does not stain cardiac or skeletal muscle; however, it does stain myofibroblasts and myoepithelial cells. This antibody could be used together with anti-muscle specific actin and myogenin in making a diagnosis of smooth muscle and skeletal muscle tumors. In most cases of rhabdomyosarcoma, this antibody yields negative results whereas anti-muscle specific actin and myogenin are positive. Leiomyosarcomas are positive only with anti-muscle specific actin and anti-smooth muscle actin and are negative with anti-myogenin.
	Immunogen	(1A4); Recombinant full-length human ACTA2 protein (ACTA2/791) (Uniprot:



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.
·	

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47699AF405

H00000059-P01-10ug	Recombinant Human alpha-Smooth Muscle Actin GST (N-Term) Protein
DVE00	VEGF [HRP]
NBP2-66429	Mouse alpha-Smooth Muscle Actin ELISA Kit (Colorimetric)
233-FB-025	FGF basic/FGF2/bFGF [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47699AF405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

