# **Product Datasheet**

## Calponin 1 Antibody (CNN1/832 + CALP) [Alexa Fluor® 750] NBP2-47758AF750

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47758AF750

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-47758AF750

#### NBP2-47758AF750

Calponin 1 Antibody (CNN1/832 + CALP) [Alexa Fluor® 750]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneCNN1/832 + CALPPreservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1264Gene ID1264Gene SymbolCNN1Specificity/SensitivityMuticle MarkerSpecificity/SensitivityMuticle MarkerSpecificity/SensitivityMuticle isoforms exist: a 34kDa form and a 29kDa form. Expression of the 29kDa form. I-calponin, is primarity restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Westro of calponin, firs and componyosin binding protein, which is thought to be involved in the regulation of smooth muscle eontraction. Calponin expression is restricted to smooth muscle eontraction. Calponin expr		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneCNN1/832 + CALPPreservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1264Gene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, 1-calponin, is primarily restricted to muscle of the urogenital antibody reacts with only the 34kDa form of calponin in extracts of culting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of unuman foreskin. Calponin is restricted to smooth muscle contraction in wascular and visceral smooth muscle. In Wester blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of unuman foreskin. Calponin is restricted to smooth muscle of unuscle of human foreskin. Calponin is restricted to smooth muscle of unuscle of human foreskin. Calponin is restricted to smooth muscle of unuscle or thread in vascular and visceral smooth muscle. In Wester blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of utilivated human foreskin. Calponin is restricted to smooth muscle colla and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneCNN1/832 + CALPPreservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1264HostMouseGene ID1264Gene SymbolCNN1Specificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, 1-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Unit Size	0.1 ml
ClonalityMonoclonalCloneCNN1/832 + CALPPreservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1264Gene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of smooth muscle during which is thought to be involved in the regulation of smooth muscle contraction. Calponin in serticted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Concentration	
CloneCNN1/832 + CALPPreservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1264HostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of cultivated human foreskin. Calponin expression is in extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityWoltpe isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of cultivated human foreskin. Calponin is a cractive with fibroblast extracts of cultivated human foreskin. Calponin is a cractive with fibroblast extracts of cultivated human foreskin. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, 1-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle. Calponin is a calmodulin, F-actin and troponyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Clone	CNN1/832 + CALP
ConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, 1-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of numan aortic medial smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 54kDa form of calponin in extracts of numan aortic medial smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 54kDa form of calponin in extracts of numan aortic medial smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 54kDa form of calponin in extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Preservative	0.05% Sodium Azide
Purity       Protein A or G purified         Buffer       50mM Sodium Borate         Product Description          Host       Mouse         Gene ID       1264         Gene Symbol       CNN1         Species       Human, Rat         Marker       Smooth Muscle Marker         Specificity/Sensitivity       Multiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, 1-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of numan aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.         Immunogen       Recombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Isotype	IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, 1-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Conjugate	Alexa Fluor 750
Product DescriptionHostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Purity	Protein A or G purified
HostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Buffer	50mM Sodium Borate
Gene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is urreactive with fibroblast extracts of cultivated human foreskin. Calponin is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Product Description	
Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Host	Mouse
SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Gene ID	1264
MarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Gene Symbol	CNN1
Specificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Species	Human, Rat
<ul> <li>molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.</li> <li>Immunogen</li> </ul>	Marker	Smooth Muscle Marker
	Specificity/Sensitivity	molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of
	Immunogen	

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-47758AF750

IC002S	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 750]
NBP1-98945-100ug	Recombinant Human Calponin 1 His Protein
1129-ER-050	ErbB2/Her2 [Unconjugated]
NBL1-09314	Calponin 1 Overexpression Lysate

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47758AF750

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

