Product Datasheet

Glypican 3 Antibody (SPM595) [Alexa Fluor® 647] NBP2-47761AF647

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47761AF647

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47761AF647



NBP2-47761AF647

Glypican 3 Antibody (SPM595) [Alexa Fluor® 647]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSPM595Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionGPC3HostMouseGene ID2719Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression of GPC3 is normal applicary carcinoma. Expression of GPC3 is normal applicary carcinoma at 50% in papillary carcinoma. Significantly higher than that of follicular adenoma. In contrast, GPC3 is nor expression and GPC3 is normal adplagent tissue. In patients with throrid cancer: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is nort expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3		-
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSPM595Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID2719Gene SymbolGPC3Specificity/SensitivityGilypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, scypression of GPC3 is dramatically enhanced in certain types of cancer, scypression of GPC3 is dramatically enhanced in certain types of ancer, scypression of GPC3 is dramatically enhanced in certain types of ancer, scypression of GPC3 is dramatically nepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, scypression of GPC3 is dramatically enhanced in certain types of ancer, scypression of GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM595 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 2719 Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcinoma. Immunogen A recombinant fragment containing amino acids 511-580 of human Glypican 3	Unit Size	0.1 ml
ClonalityMonoclonalCloneSPM595Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description4000000000000000000000000000000000000	Concentration	•
CloneSPM595Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2719Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is on depatoblastoma, with a low or undetectable expression in normal adjacent in spallary carcinoma. Expression of GPC3 is not expressed in anaplastic carcinoma. In contrast, GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2719Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. In contrast, GPC3 is not expressed in anaplastic carcinoma. In contrast, GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2719Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Clone	SPM595
ConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2719Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, melanoma, testicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2719Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Isotype	IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID2719Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Conjugate	Alexa Fluor 647
Product DescriptionHostMouseGene ID2719Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma is significantly higher than that of follicular adenoma. In contrast, GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Purity	Protein A or G purified
HostMouseGene ID2719Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Buffer	50mM Sodium Borate
Gene ID2719Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Product Description	
Gene SymbolGPC3SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Host	Mouse
SpeciesHumanMarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Gene ID	2719
MarkerHepatocellular Carcinoma MarkerSpecificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Gene Symbol	GPC3
Specificity/SensitivityGlypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma is significantly higher than that of follicular adenoma. In contrast, GPC3 is not expressed in anaplastic carcinoma.ImmunogenA recombinant fragment containing amino acids 511-580 of human Glypican 3	Species	Human
 which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma is significantly higher than that of follicular adenoma. In contrast, GPC3 is not expressed in anaplastic carcinoma. Immunogen 	Marker	Hepatocellular Carcinoma Marker
	Specificity/Sensitivity	which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma is significantly higher than that of follicular adenoma. In contrast, GPC3 is not expressed in
	Immunogen	

www.novusbio.com



Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47761AF647

IC002R	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 647]
NBP1-85226PEP	Glypican 3 Recombinant Protein Antigen
233-FB-025	FGF basic/FGF2/bFGF [Unconjugated]
2119-GP-050/CF	Glypican 3 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47761AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

