Product Datasheet

Glypican 3 Antibody (1G12 + GPC3/863) [HRP] NBP2-47763H

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47763H

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-47763H



NBP2-47763H

Product Information	
Concentration Please see the vial label for concentration. If unlisted please contact technic services. Storage Store at 4C in the dark. Clonality Monoclonal Clone 1G12 + GPC3/863 Preservative No Preservative Isotype IgG1 Kappa Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2719 Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition. GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblaston melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastor with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma and Speciarioma.	
Storage Store at 4C in the dark. Clonality Monoclonal Clone 1G12 + GPC3/863 Preservative No Preservative Isotype IgG1 Kappa Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2719 Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastor melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastor with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin (IG12); Recombinant fragment containing amino acids 511-580 of human Glypican (1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprotein)	
Clone Clone 1G12 + GPC3/863 Preservative No Preservative Isotype IgG1 Kappa Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description Host Gene ID 2719 Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBs is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastor with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin (Inc.), epotoblastor with a low or undetectable expression of GPC3 is of patients thyroid cancer, expression of GPC3 is of armatically enhanced in certain type cancers: 100% in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin (Inc.) (cal
Clone 1G12 + GPC3/863 Preservative No Preservative Isotype IgG1 Kappa Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2719 Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblaston melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastor with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain typic cancers: 100% in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin Immunogen Recombinant fragment containing amino acids 511-580 of human Glypican (1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprotein CPC3/863) (Uniprotein CPC3/863)	
Preservative Isotype IgG1 Kappa Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2719 Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastom with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcin Immunogen Recombinant fragment containing amino acids 511-580 of human Glypican (1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprotein)	
IgG1 Kappa	
Conjugate HRP Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2719 Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastor with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin (Immunogen Recombinant fragment containing amino acids 511-580 of human Glypican (1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprotes)	
Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2719 Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastom with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin (IG12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprotein P51654)	
Product Description Host Mouse Gene ID 2719 Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastom melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastor with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin (1G12); Recombinant fragment containing amino acids 511-580 of human Glypican (1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprotes)	
Product Description Host Mouse Gene ID 2719 Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastor with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain typic cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 is not expressed in anaplastic carcin follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin (1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprot P51654)	
Mouse	
Gene Symbol GPC3 Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastor melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastor with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin munogen Recombinant fragment containing amino acids 511-580 of human Glypican (1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprotential carcinoma and carcinoma carcinoma and carcinoma carcin	
Gene Symbol Species Human Marker Hepatocellular Carcinoma Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblaston with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin Recombinant fragment containing amino acids 511-580 of human Glypican (1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprotes)	
Specificity/Sensitivity Hepatocellular Carcinoma Marker	
Marker Specificity/Sensitivity Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastom melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastom with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin (1G12); Recombinant fragment containing amino acids 511-580 of human Glypican (1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprotential part of the protein (GPC3/863) (Uniprotential part of the protein (GPC3/863) (Uniprotential part of the protein (GPC3/863) (Uniprotein P51654)	
Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastor melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastor with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain typic cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin (GPC3/863) (Uniprof. P51654)	
Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by prepost-natal overgrowth and is a recessive X-linked condition.GPC3 may also found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastor melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastor with a low or undetectable expression in normal adjacent tissue. In patients thyroid cancer, expression of GPC3 is dramatically enhanced in certain type cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma was significantly higher than the follicular adenoma. In contrast, GPC 3 is not expressed in anaplastic carcin Recombinant fragment containing amino acids 511-580 of human Glypican (1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprote P51654)	
(1G12); Recombinant full-length human GPC3 protein (GPC3/863) (Uniprotein P51654)	na, ma, with es of
Product Application Details	
- Todalot Application Dotalic	
Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence	
Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence CyTOF-ready	€,
Application Notes Optimal dilution of this antibody should be experimentally determined.	
	es of at of noma.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-47763H

NBP1-43319H-0.5ml Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [HRP]

NBP1-85226PEP Glypican 3 Recombinant Protein Antigen 233-FB-025 FGF basic/FGF2/bFGF [Unconjugated]

2119-GP-050/CF Glypican 3 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47763H

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

