Product Datasheet

p57 Kip2 Antibody (SPM308) [DyLight 488] NBP2-47765G

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47765G

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47765G



NBP2-47765G

p57 Kip2 Antibody (SPM308) [DyLight 488]

stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are tripiol. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choricarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast in cuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as ar internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry. Immunofluorescence, Immunohistochemistry.		2 1
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM308 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKNIC Species Human, Mouse Species Human, Mouse Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactic with p27Kip1. p37Kip2 is a potent tight-binding inhibitor of several G1 cyclin mole (CHM) (no nuclear labeling of cyctorophoblasts and stormal cells from non-molar gestations. The histological differentiate complete moles from non-molar gestations. The histological differentiate complete moles from non-molar gestations. The histological differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clonality Monoclonal Clone SPM308 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Species Human, Mouse Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactic with p27Kip1 - p57Kip2 is a potent light-binding inhibitor of several 61 cyclin nuclear labeling of cyclotrophoblasts and stromal cells) from parial hydatidform moles, but will not differentiator of completer moles Carp. And hydropic spontaneous abortion is problematic. Most completer hydatidform moles are diploid, whereas most partial moles and hydropic spontaneous baortion. By roblematic And hydropic spontaneous baortis problematic Most complete moles from non-molar gestatio	Unit Size	0.1 ml
Clonality Monoclonal Clone SPM308 Preservative 0.05% Sodium Azide isotype IgG2b Kappa Conjugate DyLight 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKM1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 5rkDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin or uclear labeling of cytotrophoblasts and stromal cells Stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete Mydatidform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete moles and hydropic spontaneous abortion is problematic. Most complete moles and cholos. Similar moles, and hydropic islond, whereas most partial moles and hydropic islond noreas areadio islands (UTIs) demonstrate nuclear labeling i	Concentration	·
Clone SPM308 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 488 Purity Protein A or G purified Buffer S0mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactic with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells stain. The histological differentiation of complete moles form non-molar gestations. Complete moles chrom hydropic spontaneous abortion is problematic. Most complete hydatidform moles are diploid, whereas most partial moles are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervilous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as ar internal control. Immunogen Recombiant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Species Human, Mouse Species Human is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatiform mole (PHM) in which both cytotrophoblasts and stromal cells strain. The histological differentiate complete moles from nonclar gestations. Complete moles are triploid. Plody studies will identify partial moles, but will not differentiate complete moles from non-nolar gestations. Complete moles are triploid, blody studies will identify partial moles, but will not differentiate complete moles from molar gestations. Complete moles from non-loar gestations. Complete moles carry a high risk of persistent disease and choricocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblasts and stromal cells if moles are are diploid, whereas most partial moles are tabled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblasts is an	Clonality	Monoclonal
Isotype IgG2b Kappa Conjugate DyLight 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Species Human, Mouse Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactic with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete widatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblasts nucle and stromal cells are labeling with this antibody. Similar tindings apply to PHM and hydropic abortus tissues. Intervillous trophoblasti	Clone	SPM308
Conjugate DyLight 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactic with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles, and by dropic spontaneous abortion is problematic. Most complete hydatidiform moles, and by dropic aspontaneous abortion is problematic. Most complete hydatidiform moles, and hydropic abortus tisands (IVTIs) demonstrate nuclear labeling in all three entities and serve as ar internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactic with p27Kip1, p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells from partial hydatidiform moles are diploid, whereas most partial moles, and is noted and whydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partia moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriccarionma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclear labeling in all three entities and serve as ar internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Detaills Fl	Isotype	IgG2b Kappa
Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactic with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells stain. The histological differentiation of complete moles. partial moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partia moles, but will not differentiate complete moles from non-molar gestations. Complete moles from non-molar gestations. Complete moles for non-molar gestations. Complete moles for unorical ends stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as ar internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLipht (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Conjugate	DyLight 488
Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactic with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells is stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are triploid, whereas most partial moles are triploid. Ploidy studies will identify partia moles, but will not differentiate complete moles from non-molar gestations. Complete moles are triploid, ploidy studies will identify partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromat cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervinious trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as ar internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLight (R) is a trademark of Thermo Fisher Scientific I	Purity	Protein A or G purified
Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactic with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete moles, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partia moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nucle and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as ar internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. <	Buffer	50mM Sodium Borate
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactio with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, ant hydatidiform moles are diploid, whereas most partial moles from non-molar gestations. Complete moles from portula hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partia moles, but will not coliferentiate complete moles from non-molar gestations. Mouse Notes Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunohistochemistry/Immunofluorescence, Immunohis	Product Description	
Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial moles, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partia moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as ar internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunofluo	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reactic with p27Kip1, p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partia moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as ar internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunofluorescence,	Host	Mouse
Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells) stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles, but will not differentiate complete moles from non-molar gestations. Complete moles, but will not differentiate complete moles from non-molar gestations. Complete moles, but will not differentiate complete moles from non-molar gestations. Complete moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervilous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as ar internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry.Paraffin, CyTOF-ready,	Gene ID	1028
Specificity/SensitivityRecognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partia moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as ar internal control.ImmunogenRecombinant full-length human p57 Kip2 protein (Uniprot: P49918)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready,	Gene Symbol	CDKN1C
with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are 	Species	Human, Mouse
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready,	Specificity/Sensitivity	complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready,	Immunogen	Recombinant full-length human p57 Kip2 protein (Uniprot: P49918)
Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready,	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready,	Product Application Details	
	Applications	



	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47765G

NBP1-43317GMouse IgG2b Kappa Light Chain Isotype Control (MG2b) [DyLight 488]NBP1-89917PEPp57 Kip2 Recombinant Protein Antigen292-G2-050IGF-II/IGF2 [Unconjugated]NB500-106PCNA Antibody (PC10)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47765G

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

