Product Datasheet p57 Kip2 Antibody (KIP2/880) [PE] NBP2-47766PE

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47766PE

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47766PE



NBP2-47766PE

p57 Kip2 Antibody (KIP2/880) [PE]

Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monocional Clone KIP2880 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate PE Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57KD2, is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Arti-p57 has been used as an aide in identification of complete hydatiform mole (CHM) (no mole are labeling of cyctorpohoblasts and stromal cells from non-molar gestroms. The histological differentiate on the are avery tow risk. In ormal placenta, may while partial moles are tiploid, whereas most partial moles are tiploid, Pioly stutidiform mole (CHM) (no mole cell proliferation. Arti-p57 has bee	Product Information	
services. Storage Store at 4C in the dark. Clonality Monocclonal Clone KIP2/880 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate PE Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene ID 1028 Species Human, Mouse Species Human, Mouse Species Human, Mouse Species Human, Mouse Gene ID 1028 Gene ID 1028 Gene ID 1028 Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with shot hot cytorphoblasts and stromal cells stain. The histological differentiation of complete hydatidform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells is ation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblast nod storeaction withis hot chy cytotrophoblast nod storeactino	Unit Size	0.1 ml
Clonality Monoclonal Clone KIP2/880 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate PE Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1, p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin completer, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidform mole (CHM) (no nuclear labeling of cyclotrophoblasts and stromal cells) from partial hydatidform mole (PHM) in which both cytotrophoblasts and stromal cells from partial hydatidform mole (PHM) in which both cytotrophoblasts and stromal cells strom, Anno explete moles, and this antibody. Similar findings apply to PHM and hydropic aborus trophoblastic islands (I/TIs) demonstrate nuclear labeling of unrise in nor-molar gestations. Complete moles, and this antibody. Similar findings apply to PHM and hydropic aborus trophoblastic islands (I/TIs) demonstrate nuclear labeled with his antibody. Similar findings apply to PHM and hydropic caborus trophoblastic islands (I/TIs) demonstrate nuclear labeleing in all three entities a	Concentration	
Clone KIP2/880 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate PE Purity Protein A or G purified Buffer PES Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Species Human, Mouse Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells stain. The histological differentiation of complete moles, and hydropic spontaneous abortion is problematic. Most complete hydatidform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles from non-moles from non-moles from ton-moles from ton-moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriccarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar finding	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate PE Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete hydatidiform moles are diploid, whereas most partial moles, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and chorlocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands. (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control.	Clonality	Monoclonal
isotype IgG2b Kappa Conjugate PE Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Species Human, Mouse Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are tipoid. Ploidy studies will identify partial moles, but will not differentiate complete moles for non-molar gestations. Complete moles corriona, while partial moles have a very low risk of persistent disease and choropic islands (UTIs) demonstrate nuclear labeling in all three entities and serve as an internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Product Application Details <	Clone	KIP2/880
Conjugate PE Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells stain. The histological differentiation of complete moles from non-mole cells stain. The histological differentiate complete moles from non-mole cells stain. The histological differentiate complete moles from non-moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles are labeled with is antibody. Similar findings apply to PHM and hydropic aborts its uses. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Product Application Details Flow Cytometry. Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-ready.	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (PHM) (no nuclear labeling of cytotrophoblasts and stromal cells stain. The histological differentiation of complete moles from non-molar gestations. Complet moles carry a hip risk of persistent disease and choriocarcinoma, while partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles from non-molar gestations. Complete moles from and choriocarcinoma, while partial moles tare labeled with its antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Product Application Details Flow Cytometry. Immunocytochemistry/ Immunofluorescence, Immunofluorescence, Immunofluorescence,	Isotype	IgG2b Kappa
Buffer PBS Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1, p57Kip2 is a potent right-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidform mole (CHM) (no nuclear labeling of cytorophoblasts and stromal cells stain. The histological differentiation of complete moles, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidform moles are diploid, whereas most partial moles are tripiod. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar greatations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placend, horizorinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblasts manuels are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (I/TIS) demonstrate nuclear labeling in all three entities and serve as an internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) <	Conjugate	PE
Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1, p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells are main. The histological differentiate orgplete moles, partial mole, and hydropic spontaneous abortion is problematic. Most complete Hydatidiform moles are diploid, whereas most partial moles are tripioid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stormal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in Uniprot. P49918) Product Application Details Flow Cytometry, Immunocytochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunofluorescence, CyTOF-ready.	Purity	Protein A or G purified
Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1, p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complexes and thydropic spontaneous abortion is problematic. Most completes will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervilous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-ready. Recommended Dilutions Flow Cytomet	Buffer	PBS
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete moles, partial moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Product Application Detailis Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunocytochemistry-Paraffin, CyTOF-ready, Immunofluorescence, CyTOF-ready	Product Description	
Gene ID 1028 Gene Symbol CDKN1C Species Human, Mouse Specificity/Sensitivity Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are tripioid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles are a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVT1s) demonstrate nuclear labeling in all three entities and serve as an internal control. Immunogen Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene SymbolCDKN1CSpeciesHuman, MouseSpecificity/SensitivityRecognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control.ImmunogenRecombinant full-length human p57 Kip2 protein (Uniprot: P49918)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceApplicationsFlow Cytometry, Immunohistochemistry. Paraffin, CyTOF-ready, Immunofluorescence, CyTOF-ready	Host	Mouse
SpeciesHuman, MouseSpecificity/SensitivityRecognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control.ImmunogenRecombinant full-length human p57 Kip2 protein (Uniprot: P49918)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceApplicationsFlow Cytometry, Immunohistochemistry, Immunofluorescence, CyTOF-ready	Gene ID	1028
Specificity/SensitivityRecognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cyctorophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytorophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control.ImmunogenRecombinant full-length human p57 Kip2 protein (Uniprot: P49918)Product Application Details ApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Gene Symbol	CDKN1C
with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control.ImmunogenRecombinant full-length human p57 Kip2 protein (Uniprot: P49918)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunofluorescenceApplicationsFlow Cytometry, Immunohistochemistry, Immunofluorescence, Immunofluorescence, Immunohistochemistry, Immunofluorescence, CyTOF-ready	Species	Human, Mouse
Product Application Details Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready	Specificity/Sensitivity	with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma,
ApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready		findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control.
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready	Immunogen	findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control.
Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready		findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control.
Application NotesOptimal dilution of this antibody should be experimentally determined.	Product Application Details	findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control. Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready,
	Product Application Details Applications	findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control. Recombinant full-length human p57 Kip2 protein (Uniprot: P49918) Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry, Immunofluorescence,





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47766PE

NBP1-43317PE	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [PE]
NBP1-89917PEP	p57 Kip2 Recombinant Protein Antigen
292-G2-050	IGF-II/IGF2 [Unconjugated]
NB500-106	PCNA Antibody (PC10)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47766PE

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

