Product Datasheet

Golgi Glycoprotein 1/GLG1 Antibody (GLG1/970) [DyLight 755] NBP2-47794IR

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47794IR

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47794IR

NBP2-47794IR

Golgi Glycoprotein 1/GLG1 Antibody (GLG1/970) [DyLight 755]

stated on the datasheet.HostMouseGene ID2734Gene SymbolGLG1SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclor antibody is an excellent marker for human cells in xenographic model research It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneGLG1/970Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateDyLight 755PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID2734Gene SymbolGLG1SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla antibody can be used to stain the Golgi complex in cells. This monoclonal antibody use as a Golgi marker in subcelluar fractions. In produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent in tarker for human cells in kondous staten or human cells in the congraphic model research it treacts specifically with human cells in the fore one stated present and scorees and package macromolecules synthesized by the cell for exocytosis to use within the cell. Th Golgi is made up of a stack of fibrenda, research and cells in the ordigi apparatus is a organelle present in all eukaryotic cells that forms a part of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within th	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneGLG1/970Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateDyLight 755PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID2734Gene SymbolGLG1SpeciesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and 16 E-selectin. (Gel-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. (He and malgrant cells. This monoclonal antibody racing and 16 Cyserio GLG1 repeats. This monoclonal antibody cell apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi zone in normal and malgrant cells. This monoclonal antibody is an excellent marker for human cells in the cell of resceryois or use within the cell. The Golgi apparatus is to process and package macromolecules synthesized by the cell of resceryois or use within the cell. The Golgi apparatus is to process and package macromolecules synthesized by the cell of resceryois or use within the cell. To Golgi is macke up of a stack of flattened, membrane-bound sack known as cistermae, with three functional regions: the cell so the more pronounced and numerous in cells that make and secrete many substances such	Unit Size	0.1 ml
Clonality Monoclonal Clone GLG1/970 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 755 Purity Protein A or G purified Buffer S0mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated on uter of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 2734 Gene Symbol GLG1 Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. I his monoclonal antibody can be used to stain the Golgi apparatus is an organelle present in all eukaryotic cells that forms apart of the endomembrane system. The primary function of the Golgi apparatus is to process and package. This monoclonal antibody is an excellent marker for human cells. The songraphic model research for accoptic size used to produce shown as cistemae, with three functional regions: the cis face, medial region and trans face. Each region consists of fatten	Concentration	•
Cione GLG1/970 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 755 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 2734 Gene Symbol GLG1 Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating and can be used as a Golgi ranker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi apparatus is to process and package maccronlecules synthesized by the cell for exocytosis or use within the cell. The primary function of the Golgi apparatus is to process and package maccronlecules passing though them, depending on where they are destined unmeronlecules passing though them, depending on wher	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 755 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 2734 Gene Symbol GLG1 Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin. It contains stalic acid residues and reading or multipaparatus is an organelle growth factor shale used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus arring pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus expresent in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi aparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi aparatus is an organelle macromolecules synt	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateDyLight 755PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated oulme of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID2734Gene SymbolGLG1SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neurophils). Fucosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GL4T repeats. This monoclonal antibody can be used to stain the Golgi complex in cells on tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is the dialare gion and trans cister. Each region consists of various enzymes that selectively modify the macromolecules systems do various enzymes that selectively modify the macromolecules system side so thave budded off of the Golgi represent surrounding the main cisternae. The Golgi tends to be more pronounced and numeromlecules apherical v	Clone	GLG1/970
ConjugateDyLight 755PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID2734Gene SymbolGLG1SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It reactions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monocloral antibody is an excellent marker for human and malignant cells. This monocloral antibody is an excellent marker for binding of the essential for binding to E-selectin. It reacts specifically with human cells in xenographic model research threacts specifically with human cells in xenographic model research threacts specifically with human cells in xenographic model research threacts and eave of a stack of flattened, membrane-bound sacs known as cistermae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi apparatus is an organelle present in all eukaryotic cells that have budded off of the Golgi apparatus is an organelle present in all eukaryotic vesides that have budded off	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 2734 Gene Symbol GLG1 Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. If the Golgi zone in normal and malignant cells. This monoclonal antibody can be used to stain the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research the reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research is the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cistemare, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the marcomolecules spassing though them,	Isotype	IgG1 Kappa
Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 2734 Gene Symbol GLG1 Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is an organelle present with tree clus specifically with human cells. The Golgi apparatus is an organelle present with the cell matice on the didapaparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, metharens-bound sack Known as cistermae, with three functional regions: the cis face, medial reg	Conjugate	DyLight 755
Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 2734 Gene Symbol GLG1 Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating to the binding of neutrophils). Fuccosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cells or this an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B cells.	Purity	Protein A or G purified
DescriptionThis conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID2734Gene SymbolGLG1SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialia caid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi and there donembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi and the asc have budded off of the Golgi are present stace. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined reside. Several spherical vesices that have budded off of the Golgi are present surrounding the main cisternae. The Golgi tost be more pronounced and numerous in cells that make and secrete many substances such as plasma B cells. </th <th>Buffer</th> <th>50mM Sodium Borate</th>	Buffer	50mM Sodium Borate
volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.HostMouseGene ID2734Gene SymbolGLG1SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sailic acid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosics or use within the cell. Th Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B 	Product Description	
Gene ID2734Gene SymbolGLG1SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclor antibody is an excellent marker for human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B cells.ImmunogenGolgi fraction from human liver cells	Description	volume of this product. The volume will be greater than or equal to the unit size
Gene SymbolGLG1SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclor antibody is an excellent marker for human cells in xenographic model research It reacts specifically with human cells in xenographic model research It reacts specifically with human cells in the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined reside. Several spherical vesicles that have budded off the Golgi are present surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B cells.ImmunogenGolgi fraction from human liver cells	Host	Mouse
SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin.Specificity/SensitivityThis monoclonal antibody recognizes and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclor antibody is an excellent marker for human cells in xenographic model research It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined reside. Several spherical vesicles that have budded off of the Golgi ap resent surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B cells.ImmunogenGolgi fraction from human liver cells	Gene ID	2734
Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclor antibody is an excellent marker for human cells in xenographic model research It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B cells.ImmunogenGolgi fraction from human liver cells	Gene Symbol	GLG1
MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclor antibody is an excellent marker for human cells in xenographic model research It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B cells.ImmunogenGolgi fraction from human liver cells	Species	Human, Mouse (Negative), Rat (Negative)
Specificity/SensitivityThis monoclonal antibody recognizes a protein of 134kDa, which binds fibrobla growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclor antibody is an excellent marker for human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B cells.ImmunogenGolgi fraction from human liver cells	Reactivity Notes	Does not react with Mouse or Rat.
growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GLG1 repeats. This monoclonal antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffus staining pattern of the Golgi zone in normal and malignant cells. This monoclor antibody is an excellent marker for human cells in xenographic model research It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. Th Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B cells.ImmunogenGolgi fraction from human liver cells	Marker	Marker for Human Cells
	Specificity/Sensitivity	antibody can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B
Notes		Golgi fraction from human liver cells
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.



Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.



technical@novusbio.com



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47794IR

NBP1-43319IR-0.5ml	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 755]
NBP1-90636PEP	Golgi Glycoprotein 1/GLG1 Recombinant Protein Antigen
233-FB-025	FGF basic/FGF2/bFGF [Unconjugated]
BBA16	E-Selectin/CD62E Antibody (BBIG-E4) [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47794IR

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

