# **Product Datasheet**

# Fc gamma RIIA/CD32a Antibody (7.3) [Janelia Fluor® 549] NBP2-47830JF549

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47830JF549

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-47830JF549



# NBP2-47830JF549

Fc gamma RIIA/CD32a Antibody (7.3) [Janelia Fluor® 549]

Fc gamma RIIA/CD32a Antibody (7.3) [Janelia Fluor® 549]	
0.1 ml	
Please see the vial label for concentration. If unlisted please contact technical services.	
Store at 4C in the dark.	
Monoclonal	
7.3	
0.05% Sodium Azide	
IgG1 Kappa	
Janelia Fluor 549	
Protein A or G purified	
50mM Sodium Borate	
Mouse	
2212	
FCGR2A	
Human	
This monoclonal antibody reacts with a CD32 (FcgRII) epitope (cluster-4). It displays a stronger reaction with Daudi than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab'2 fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoforms are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 and IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms. In addition, the IIb isoform is expressed by B cells, and the IIa isoform by platelets, granulocytes and, weakly, by B cells. NK cells and neutrophils express Isoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexes.	
K562 and FcgRII+L cells	
Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus	
Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready, Immunofluorescence	
Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-ready	
Optimal dilution of this antibody should be experimentally determined.	





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## Products Related to NBP2-47830JF549

NBP1-84589PEP Fc gamma RIIA/CD32a Recombinant Protein Antigen

210-TA-005 TNF-alpha [Unconjugated]

1330-CD-050 Fc gamma RIIA/CD32a [Unconjugated]

M6000B-1 IL-6 [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47830JF549

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

