Product Datasheet

Fc gamma RIIA/CD32a Antibody (8.7) [Allophycocyanin/Cy7] NBP2-47831APCCY7

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47831APCCY7

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47831APCCY7

NBP2-47831APCCY7

Fc gamma RIIA/CD32a Antibody (8.7) [Allophycocyanin/Cy7]

services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone 8.7 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Allophycocyanin/Cy7 Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da transmembrane glycoprotein in that mediates several functions including phagocytosis, cytoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoform are generated via alternative mRNA splicing, i.e., 114, 1182, IID3 and the serversa all isoform is expressed by B cells, and neutrophils express lisoform in are director in the region of monomeric Ig6 but more storing is to add end by B cells, and neutrophils express lisoform is corres and isoform is expressed by B cells, and neutrophils express lisoform is corres and isoform is expressed by B cells, and neutrophils express lisoform is corres and endothelial cells Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-read Immunofluorescence.		
Concentration Please see the vial label for concentration. If unlisted please contact technic services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone 8.7 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate AllophycocyanirCy7 Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 2212 Gene Symbol FCGR2A Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 8.26 and the epitope overlaps with that of monoclonal antibody 7.30 (kupsys a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complex binding. CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 8.26 and the epitope ourlaps with that of monoclonal antibody 7.30 (kupsys a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complex binding. CD32 (FcgRII) a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and a teast 6 isoforn are generated via alternative mRNA splicing, i.e., IIa1, IIb1, IIb2, IIb3 a ticoform by platelets, granulocytes and, weakly, by B cells.	Product Information	
services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone 8.7 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Allophycocyanin/Cy7 Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da transmembrane glycoprotein in that mediates several functions including phagocytosis, cytoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoform are generated via altemative mRNA splicitorin is expressed by B cells, and IL isoform is expressed and inform is expressed by B cells, and IL isoform is expressed by B cells, and IL is	Unit Size	0.1 ml
Clonality Monocional Clone 8.7 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Allophycocyanin/Cy7 Purity Protein A or G purified Buffer PBS Product Description Host Host Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 8.26 and the epitope overlaps with that of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U37 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complexe binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoform are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 a IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoform is expressed by B cells, and IIa isoform by platelets, granulocytes and, weakly, b B cells. NK cells and neutrophils express Isoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe Immunofluorescence, CyTOF-ready Product Application Details Flow Cytometry, Immunocytochemistry/ Immu	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Clone 8.7 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Allophycocyanin/Cy7 Purity Protein A or G purified Buffer PBS Product Description Product Description Host Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody s.26 and the epitope overlaps with that of monoclonal antibody reacts with a CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoform are generated via altemative mRNA splicing, i.e., IIa1, IIB2, IIb2, IIb3 a IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoform is expressed by B cells, and IIa isoform by platelets, granulocytes and, weakly, by B cells. NK cells and neutrophils express Isoform II.c. CD32 binds weakly to the Fc region of monomeric Ig6 but more strongly to IgG aggregates and immune complexe Immunofluorescence, CyTOF-read/Immunofluorescence, CyTOF-read/Immunofluorescence, CyTOF-read/Immunofluorescence, CyTOF-read/Immunofluorescence, CyTOF-read/Immunofluorescence, CyTOF-read/Immunofluorescence, CyTOF-read/Immunofluorescence, CyTOF-read/Immunofluorescence, CyTOF-read/Immunofluorescence, CyTOF-read/Immunofluorescence/Immunofluorescence/Immunofluorescence/Immu	Storage	Store at 4C in the dark. Do not freeze.
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Allophycocyanin/Cy7 Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Species Species	Clonality	Monoclonal
Isotype IgG1 Kappa Conjugate Allophycocyanin/Cy7 Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domina collection of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with that of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction sincluding transmembrane glycoprotein that mediates several functions including hagocytosis, cytotoxidity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoform are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 a IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express al isoforms. In addition, the IIb isoform is expressed by B cells. NK cells and neutrophils express Isoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready Application Notes Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Sortex or externe temperature fluctuations. These can lead to inreversible degrada or decoupling. When staining intracellular targets, specific attention to t	Clone	8.7
Conjugate Allophycocyanin/Cy7 Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 8.26 and the epitope overlaps with that of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIs. Its Fab fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicty, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoforn are generated via alternative mRNA splicing, i.e., Ital, Ita2, Itb1, Itb2, Itb3 a Itb. Monocytes/macrophages, placental tropholasts and endothelial cells express all isoforms is expressed by B cells, and Ita isoform by platelets, granulocytes and, weakly, by B cells. NK Cells and neutrophils express loform Itc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe K562 and FcgRII+L cells Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to inreversible degrada or decoupling. When staini	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer PBS Product Description Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FogRII) epitope distinct from defined by monoclonal antibody 8.26 and the epitope overlaps with that of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIa. Its Fab fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoform are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 a lic. Monocytes/macrophages, placental tropholtasts and endothelial cells express al to be cells, and neutrophils express loform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe for monomeric IgG but more strongly to IgG aggregates and immune complexe Immunofluorescence, CyTOF-ready Application Notes Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, Ortor, Immunofluorescence, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid proinged exposure to or extreme temperature fluctuations. These can lead to irreversible degrado or decoupling. When staining intracelluar tagets, specific	Isotype	IgG1 Kappa
Buffer PBS Product Description Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from odefined by monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 a IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms in addition, the IIb isoform is expressed by B cells, and IIa isoform by platelets, granulocytes and, weakly, by B cells. NK cells and neutrophils express Isoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe Immunogen K562 and FcgRII+L cells Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fi and permeabilization steps in your flow protocol may be required. Please co our technica	Conjugate	Allophycocyanin/Cy7
Product Description Host Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 8.26 and the epitope overlaps with that of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis. cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoform are generated via alternative mRNA splicing, i.e., Ila1, Ila2, Ilb1, Ilb2, Ilb3 a Ilc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms. In addition, the Ilb isoform is expressed by B cells, and Ila isoform by platelets, granulocytes and, weakly, by B cells. NK cells and neutrophils express Isoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe Immunofluorescence, CyTOF-ready Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-ready Application Notes Optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to ineversible degrada or decoupling. When staining intracellular targets, specific attention to the fi and permeabilization steps in your flow protocol may be required. Please co optimal results using our Tandem dyes, please avoid prolong	Purity	Protein A or G purified
Host Mouse Gene ID 2212 Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 8.26 and the epitope overlaps with that of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoform are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 a IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms. In addition, the IIb isoform is expressed by B cells, NK cells and neutrophils express Isoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe Immunofluorescence, CyTOF-ready Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready Application Notes Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fin and permeabilization steps in your flow pr	Buffer	PBS
Gene ID 2212 Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 8.26 and the epitope overlaps with that of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoforn are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 a IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms. In addition, the IIb isoform is expressed by B cells, and IIa isoform by platelets, granulocytes and, weakly, by B cells. NK cells and neutrophils express Isoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe Immunofluorescence Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready Immunofluorescence Application Notes Flow Cytometry, Immunocytochemistry/ Immunofluorescence, immunofluorescence, CyTOF-ready Immunofluorescence, CyTOF	Product Description	
Gene Symbol FCGR2A Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 8.26 and the epitope overlaps with that of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoform are generated via alternative mRNA splicing, i.e., Ila1, Ila2, Ilb1, Ilb2, Ilb3 a Ilc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms. In addition, the Ilb isoform is expressed by B cells, NK cells and neutrophils express lsoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe Immunofluorescence, CyTOF-ready Application Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready Immunofluorescence, CyTOF-ready Immunofluorescence, CyTOF-ready Immunofluorescence, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the finand permeabilization steps in your flow protocol may be required. Please co our technical support team at technical envolution.	Host	Mouse
Species Human Specificity/Sensitivity This monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 8.26 and the epitope overlaps with that of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoform are generated via alternative mRNA splicing, i.e., IIa1, IIb2, IIb1, IIb2, IIb3 a IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms. In addition, the IIb isoform is expressed by B cells, and neutrophils express Isoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe Immunofluorescence Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready Immunofluorescence, Immunofluorescence, CyTOF-ready Imm	Gene ID	2212
Specificity/SensitivityThis monoclonal antibody reacts with a CD32 (FcgRII) epitope distinct from defined by monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complex binding. CD32 (FcRII) is a type 1 	Gene Symbol	FCGR2A
defined by monoclonal antibody 8.26 and the epitope overlaps with that of monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Da than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoforn are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 a IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms. In addition, the IIb isoform is expressed by B cells, and IIa isoform by platelets, granulocytes and, weakly, by B cells. NK cells and neutrophils express Isoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexe Immunogen Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention the fit and permeabilization steps in your flow protocol may be required. Please co our technical support team at technical@novusbio.com if you have any	Species	Human
Product Application Details Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready Immunofluorescence Recommended Dilutions Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-ready Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please co our technical support team at technical@novusbio.com if you have any	Specificity/Sensitivity	monoclonal antibody 7.30 (cluster 4). It displays a stronger reaction with Daudi than with U937 cells. The epitope is located in domain 2 of FcgRIIa. Its Fab'2 fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoforms are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 and IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms. In addition, the IIb isoform is expressed by B cells, and the IIa isoform by platelets, granulocytes and, weakly, by B cells. NK cells and
ApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-readyRecommended DilutionsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please co our technical support team at technical@novusbio.com if you have any	Immunogen	K562 and FcgRII+L cells
ImmunofluorescenceRecommended DilutionsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, CyTOF-readyApplication NotesOptimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please co our technical support team at technical@novusbio.com if you have any	Product Application Details	
Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please co our technical support team at technical@novusbio.com if you have any	Applications	
optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please co our technical support team at technical@novusbio.com if you have any		
94004010.	Application Notes	optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47831APCCY7

NBP1-84589PEP	Fc gamma RIIA/CD32a Recombinant Protein Antigen
210-TA-005	TNF-alpha [Unconjugated]
1330-CD-050	Fc gamma RIIA/CD32a [Unconjugated]
M6000B-1	IL-6 [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47831APCCY7

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

