Product Datasheet

HSP60 Antibody (HSPD1/780) - Azide and BSA Free NBP2-47843-0.1mg

Unit Size: 0.1 mg

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47843

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47843



NBP2-47843-0.1mg


HSP60 Antibody (HSPD1/780) - Azide and BSA Free

0.1 mg	
1.0 mg/ml	
Store at -20 to -80C. Avoid freeze-thaw cycles.	
Monoclonal	
HSPD1/780	
No Preservative	
IgG1 Kappa	
Protein A or G purified	
10 mM PBS	
60 kDa	
Product Description	
 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA. Also available at 200 ug/ml WITH BSA & azide (NBP2-44758). Antibody with azide - store at 2 to 8C. 	
Mouse	
3329	
HSPD1	
Human, Mouse, Rat, Porcine, Bovine, Canine, Chicken, Hamster, Monkey, Rabbit, Sheep	
Mitochondrial Marker	
Recognizes a 60kDa protein, identified as the heat shock protein 60 (hsp60). A wide variety of environmental and pathophysiological stressful conditions trigger the synthesis of a family of proteins known as heat shock proteins (hsps). Hsp60 is a potential antigen in a number of autoimmune diseases. In human arthritis and in experimentally induced arthritis in animals, disease development coincides with the development of immune reactivity directed against not only bacterial hsp60, but also against its mammalian homolog.	
Recombinant human HSP60 protein (Uniprot: P10809)	
Product Application Details	
Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array, CyTOF- ready, Immunofluorescence	
Western Blot 0.25 - 0.5 ug/ml, Flow Cytometry 0.5 - 1 ug/million cells in 0.1 ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, Protein Array, Immunofluorescence 0.5 - 1.0 ug/ml, CyTOF-ready	
Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.	



Images

Western Blot: HSP60 Antibody (HSPD1/780) - Azide and BSA Free [NBP2-47843] - Western Blot analysis of HeLa cell lysate using HSP60 Antibody (HSPD1/780)





Immunocytochemistry/Immunofluorescence: HSP60 Antibody (HSPD1/780) - Azide and BSA Free [NBP2-47843] - Immunofluorescent staining of PFA-fixed MCF-7 cells. HSP60 Antibody (HSPD1/780) followed by goat anti-Mouse IgG-CF488 (Green). Nuclei are labeled with Red Dot (red).


Immunohistochemistry-Paraffin: HSP60 Antibody (HSPD1/780) - Azide and BSA Free [NBP2-47843] - Human Renal Cell Carcinoma stained with HSP60 Monoclonal Antibody (HSPD1/780)

Immunocytochemistry/Immunofluorescence: HSP60 Antibody (HSPD1/780) - Azide and BSA Free [NBP2-47843] - Immunofluorescent staining of MeOH-fixed HeLa cells. HSP60 Antibody (HSPD1/780) followed by goat anti-Mouse IgG-CF488 (green). Nuclei are labeled with Red Dot (red).

www.novusbio.com










Immunohistochemistry-Paraffin: HSP60 Antibody (HSPD1/780) - Azide and BSA Free [NBP2-47843] - Human Bladder Carcinoma stained with HSP60 Monoclonal Antibody (HSPD1/780)

Immunohistochemistry-Paraffin: HSP60 Antibody (HSPD1/780) - Azide and BSA Free [NBP2-47843] - Human Lung Carcinoma stained with HSP60 Monoclonal Antibody (HSPD1/780)

Protein Array: HSP60 Antibody (HSPD1/780) - Azide and BSA Free [NBP2-47843] - Analysis of Protein Array containing more than 19,000 full-length human proteins using HSP60 Antibody (HSPD1/780) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5.







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47843-0.1mg

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP1-77397PEP	HSP60 Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47843

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

