Product Datasheet

Androgen R/NR3C4 Antibody (DHTR/882) [DyLight 680] NBP2-47856FR

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47856FR

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47856FR



NBP2-47856FR

Androgen R/NR3C4 Antibody (DHTR/882) [DyLight 680]

District Concentration	
Concentration Please see the vial label for concentration. If unlisted please contact tech services. Storage Store at 4C in the dark. Clonality Monoclonal Clone DHTR/882 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer SomM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the s volume of this product. The volume will be greater than or equal to the ur stated on the datasheet. Host Mouse Gene ID 367 Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade idifferentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a m hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been us clinically in differentiating morpheaform basal cell carcinoma (mBCC) frodesmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibs superb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
Storage Store at 4C in the dark. Clonality Monoclonal Clone DHTR/882 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the s volume of this product. The volume will be greater than or equal to the ur stated on the datasheet. Host Mouse Gene ID 367 Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or gluccorticiol receptors. The expression of AR is reportedly inversely correlated with histologic grade id differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors show higher expression than the poorly differentiated prostate cancer, AR has been proposed, as a m hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been use clinically in differentiating morpheaform basal cell carcinoma (mBCC) for desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibo superb for staining of formalin/paraffin tissues. Immunogen	
Clone DHTR/882 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the s volume of this product. The volume will be greater than or equal to the ur stated on the datasheet. Host Mouse Gene ID 367 Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with sull length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade id differentiated tumors. In prostate cancer, AR has been proposed, as a nommone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been use clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibos superb for staining of formalin/paraffin tissues. Immunogen	nical
Clone DHTR/882 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the s volume of this product. The volume will be greater than or equal to the ur stated on the datasheet. Host Mouse Gene ID 367 Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade if differentiated tumors. In prostate cancer, AR has been proposed, as a m hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been us clinically in differentiating morpheaform basal cell carcinoma (mBCC) fro desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibos uperb for staining of formalin/paraffin tissues. Immunogen	
Preservative IgG1 Kappa IgG1 Kappa IgG1 Kappa DyLight 680 Purity Protein A or G purified SomM Sodium Borate	
IgG1 Kappa	
Conjugate Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the s volume of this product. The volume will be greater than or equal to the unstated on the datasheet. Host Mouse Gene ID 367 Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade id differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a m hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) frod desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibous upper for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the s volume of this product. The volume will be greater than or equal to the ur stated on the datasheet. Host Mouse Gene ID 367 Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade in differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a m hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) frodesmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibous usperb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
Product Description This conjugate is made on demand. Actual recovery may vary from the s volume of this product. The volume will be greater than or equal to the unstated on the datasheet. Host Mouse Gene ID 367 Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade in differentiated tumors. In prostate cancer, AR has been proposed, as a mention hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibous per bor staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
Product Description This conjugate is made on demand. Actual recovery may vary from the s volume of this product. The volume will be greater than or equal to the unstated on the datasheet. Host Mouse Gene ID 367 Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade id differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a m hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been us clinically in differentiating morpheaform basal cell carcinoma (mBCC) frodesmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibos superb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
This conjugate is made on demand. Actual recovery may vary from the s volume of this product. The volume will be greater than or equal to the ur stated on the datasheet. Mouse Gene ID 367 Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade id differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a m hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been use clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibos superb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
volume of this product. The volume will be greater than or equal to the ur stated on the datasheet. Mouse Gene ID 367 Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade in differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a machormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibous uperb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade in differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a man hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) from the description of the skin. This monoclonal antibodistic trichoepithelioma (DTE) in the skin.	
Gene Symbol AR Species Human, Mouse (Negative) Reactivity Notes Does not react with Mouse. Marker Marker of Androgen Dependence Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade in differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a man hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibot superb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
Species Human, Mouse (Negative) Does not react with Mouse. Marker Marker of Androgen Dependence Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade in differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a man hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) frow desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibous superb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
Reactivity Notes Marker Marker of Androgen Dependence Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade in differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a mention hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) from the desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibod superb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
Marker Marker of Androgen Dependence Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade is differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a monomone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibous superb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
Recognizes a protein of 110kDa, which is identified as androgen receptor reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade is differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a m hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibodish superb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
reacts with full length, and the newly described A form of the receptor. It cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade is differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a m hormone-responsiveness and thus it may be useful in identifying patients benefit from anti-androgen therapy. Anti-androgen receptor has been used clinically in differentiating morpheaform basal cell carcinoma (mBCC) frow desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibous superb for staining of formalin/paraffin tissues. Immunogen Human Androgen R/NR3C4 recombinant protein fragment; aa151-350 (E	
	does not e. well arker of likely to eful m
sequence is proprietary) (Onliprot. F 10273)	xact
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subside	aries.
Product Application Details	
Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence	
Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence CyTOF-ready	ice,



Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-47856FR

NBP1-43319FR-0.5ml Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 680]
H00000367-Q01-10ug Recombinant Human Androgen R/NR3C4 GST (N-Term) Protein

236-EG-200 EGF [Unconjugated]

NBP2-69854 Human Androgen R/NR3C4 ELISA Kit (Chemiluminescence)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47856FR

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

